

# STANLEY–REISNER RINGS AND TORUS ACTIONS

TARAS PANOV

**ABSTRACT.** We review a class of problems on the borders of topology of torus actions, commutative homological algebra and combinatorial geometry, which is currently being investigated by Victor Buchstaber and the author. The text builds on the lectures delivered on the transformation group courses in Osaka City University and Universitat Autònoma de Barcelona. We start with discussing several well-known results and problems on combinatorial geometry of polytopes and simplicial complexes, and then move gradually towards investigating the combinatorial structures associated with spaces acted on by the torus. Parallely, we set up the required commutative algebra apparatus, including Cohen–Macaulay/Gorenstein rings and Stanley–Reisner face rings of simplicial complexes.

## 1. COMBINATORIAL MOTIVATIONS: $f$ -VECTORS

Here we review some well-known combinatorial results and problems concerning the number of faces of simplicial polytopes and complexes.

Let  $K^{n-1}$  be an arbitrary  $(n-1)$ -dimensional simplicial complex on the vertex set  $[m] = \{1, \dots, m\}$ . The  $f$ -vector of  $K$  is defined as  $\mathbf{f}(K) = (f_0, \dots, f_{n-1})$  where  $f_i$  is the number of  $i$ -dimensional simplices of  $K$ . We also put  $f_{-1}(K) = 1$ . The equivalent information is contained in the  $h$ -vector  $\mathbf{h}(K) = (h_0, \dots, h_n)$ , defined from the equation

$$(1.1) \quad h_0 t^n + \dots + h_{n-1} t + h_n = (t-1)^n + f_0(t-1)^{n-1} + \dots + f_{n-1}.$$

Convex geometry provides an important class of simplicial complexes, the boundaries of *simplicial polytopes*. Given a simplicial polytope  $Q$ , we define its  $f$ - and  $h$ -vector as those of its boundary complex. The famous theorem of Billera–Lee and Stanley (proved in 1980) gives a complete characterisation of  $f$ -vectors of simplicial polytopes.

For any two positive integers  $a, i$  there exists a unique *binomial  $i$ -expansion* of  $a$  of the form

$$a = \binom{a_i}{i} + \binom{a_{i-1}}{i-1} + \dots + \binom{a_j}{j},$$

where  $a_i > a_{i-1} > \dots > a_j \geq j \geq 1$ . Define

$$a^{(i)} = \binom{a_i+1}{i+1} + \binom{a_{i-1}+1}{i} + \dots + \binom{a_j+1}{j+1}, \quad 0^{(i)} = 0.$$

---

The author is grateful to Mikiya Masuda of Osaka City University and Natàlia Castellana of Universitat Autònoma de Barcelona for the insight gained from numerous discussions and great hospitality. The work was partially supported by the Russian Foundation for Basic Research, grant no. 01-01-00546. Preprint id. ITEP-TH-91/02.

**Example 1.2.** For any  $a > 0$  we have  $a^{(1)} = \binom{a+1}{2}$ . Let  $a = 28$ ,  $i = 4$ . The corresponding binomial expansion is

$$28 = \binom{6}{4} + \binom{5}{3} + \binom{3}{2}.$$

Hence,

$$28^{(4)} = \binom{7}{5} + \binom{6}{4} + \binom{4}{3} = 40.$$

**Theorem 1.3** (Billera–Lee, Stanley). *An integer sequence  $(f_0, f_1, \dots, f_{n-1})$  is the  $f$ -vector of a simplicial  $n$ -polytope if and only if the corresponding sequence  $(h_0, \dots, h_n)$  determined by (1.1) satisfies the following three conditions:*

- (a)  $h_i = h_{n-i}$ ,  $i = 0, \dots, n$  (the Dehn–Sommerville equations);
- (b)  $h_0 \leq h_1 \leq \dots \leq h_{\lfloor \frac{n}{2} \rfloor}$ ;
- (c)  $h_0 = 1$ ,  $h_{i+1} - h_i \leq (h_i - h_{i-1})^{(i)}$ ,  $i = 1, \dots, \lfloor \frac{n}{2} \rfloor - 1$ .

**Example 1.4.** 1. The first inequality  $h_0 \leq h_1$  from part (b) of  $g$ -theorem is equivalent to  $f_0 = m \geq n + 1$ . This just expresses the fact that it takes at least  $n + 1$  hyperplanes to bound a polytope in  $\mathbb{R}^n$ .

2. The first inequality from part (c) is equivalent to the upper bound

$$f_1 \leq \binom{f_0}{2},$$

which says that any two vertices are joined by at most one edge.

3. The second inequality  $h_1 \leq h_2$  (for  $n \geq 4$ ) from part (b) is equivalent to the lower bound

$$f_1 \geq n f_0 - \binom{n+1}{2}.$$

**Exercise 1.5.** Deduce the following “ $f$ -vector form” of the Dehn–Sommerville equations

$$(1.6) \quad f_{k-1} = \sum_{j=k}^n (-1)^{n-j} \binom{j}{k} f_{j-1}, \quad k = 0, 1, \dots, n.$$

An integral sequence  $(k_0, k_1, \dots, k_r)$  satisfying  $k_0 = 1$  and  $0 \leq k_{i+1} \leq k_i^{(i)}$  for  $i = 1, \dots, r-1$  is called an  $M$ -vector (after M. Macaulay). One may introduce the  $g$ -vector  $\mathbf{g}(K^{n-1}) = (g_0, g_1, \dots, g_{\lfloor \frac{n}{2} \rfloor})$  of simplicial complex  $K^{n-1}$  by  $g_0 = 1$ ,  $g_i = h_i - h_{i-1}$ ,  $i > 0$ . Then conditions (b) and (c) from the  $g$ -theorem are equivalent to that the  $g$ -vector of a simplicial  $n$ -polytope is an  $M$ -vector. The notion of  $M$ -vector arises in the following theorem of commutative algebra.

**Theorem 1.7** (Macaulay). *An integral sequence  $(k_0, k_1, \dots, k_r)$  is an  $M$ -vector if and only if*

$$k_i = \dim_{\mathbf{k}} A^{2i}, \quad i = 1, \dots, r.$$

*for some commutative graded connected  $\mathbf{k}$ -algebra  $A = A^0 \oplus A^2 \oplus \dots \oplus A^{2r}$  generated by degree-two elements.*

We sketch the proof of the necessity part of Theorem 1.3 in the end of section 4.1. Below we review two more famous results concerning the number of faces, *Upper Bound* and *Lower Bound theorems* for simplicial polytopes. Both can be deduced from Theorem 1.3.

A simplicial polytope  $Q$  is called  $k$ -neighbourly if any  $k$  vertices span a face.

**Exercise 1.8.** Using the Dehn–Sommerville equations, show that if  $Q$  is a  $k$ -neighbourly simplicial  $n$ -polytope and  $k > \lfloor \frac{n}{2} \rfloor$ , then  $Q$  is an  $n$ -simplex.

A  $\lfloor \frac{n}{2} \rfloor$ -neighbourly simplicial  $n$ -polytope is called *neighbourly*. An example of a neighbourly  $n$ -polytope with arbitrary number of vertices is provided by the *cyclic polytope*  $C^n(m)$  defined as the convex hull of any  $m$  distinct points  $\mathbf{x}(t_i)$ ,  $t_1 < t_2 < \dots < t_m$ , on the *moment curve*

$$\mathbf{x}: \mathbb{R} \longrightarrow \mathbb{R}^n, \quad t \mapsto \mathbf{x}(t) = (t, t^2, \dots, t^n) \in \mathbb{R}^n.$$

**Exercise 1.9.** Prove that the  $C^n(m)$  is indeed a neighbourly simplicial polytope, and that its combinatorial type is independent on a choice of points on the moment curve.

**Theorem 1.10** (UBT for simplicial polytopes). *The number of  $i$ -faces of arbitrary simplicial  $n$ -polytope  $Q$  does not exceed the number of  $i$ -faces of any neighbourly  $n$ -polytope with the same number of vertices. That is, if  $f_0(Q) = m$ , then*

$$f_i(Q) \leq f_i(C^n(m)) \quad \text{for } i = 1, \dots, n-1.$$

*The equality above holds if and only if  $Q$  is a neighbourly polytope.*

The Upper Bound theorem was conjectured by Motzkin in 1957 and proved by P. McMullen in 1970.

Note that, since  $C^n(m)$  is neighborly,

$$f_i(C^n(m)) = \binom{m}{i+1} \quad \text{for } i = 0, \dots, \lfloor \frac{n}{2} \rfloor - 1.$$

Due to the Dehn–Sommerville equations this determines the full  $f$ -vector of  $C^n(m)$ .

**Exercise 1.11.** Calculate the  $f$ -vector of a neighbourly simplicial  $n$ -polytope.

**Exercise 1.12.** Prove that the UBT inequalities are equivalent to the following inequalities for the  $h$ -vector:

$$h_i(Q) \leq \binom{m-n+i-1}{i}, \quad i = 0, \dots, \lfloor \frac{n}{2} \rfloor.$$

(here  $m = f_0$  is the number of vertices). This was the key observation in McMullen’s proof of the UBT and its subsequent generalisation to arbitrary simplicial spheres due to Stanley. We will return to this argument later.

A simplicial  $n$ -polytope  $Q$  is called *stacked* if there is a sequence  $Q_0, Q_1, \dots, Q_k = Q$  of  $n$ -polytopes such that  $Q_0$  is an  $n$ -simplex and  $Q_{i+1}$  is obtained from  $Q_i$  by adding a pyramid over some facet of  $Q_i$ . In the combinatorial language, stacked polytopes are those obtained from a simplex by applying several subsequent stellar subdivisions of facets.

**Theorem 1.13** (LBT for simplicial polytopes). *For any simplicial  $n$ -polytope  $Q$  ( $n \geq 3$ ) with  $m = f_0$  vertices the following lower bounds hold*

$$\begin{aligned} f_i(Q) &\geq \binom{n}{i} f_0 - \binom{n+1}{i+1} i \quad \text{for } i = 1, \dots, n-2; \\ f_{n-1}(Q) &\geq (n-1) f_0 - (n+1)(n-2). \end{aligned}$$

*The equality is achieved if and only if  $Q$  is a stacked polytope.*

Note that the first inequality above is equivalent to the inequality  $h_1 \leq h_2$  from Theorem 1.3.

The boundary of a simplicial  $n$ -polytope is a simplicial subdivision of  $(n-1)$ -sphere. However, not every triangulation of sphere is combinatorially equivalent to a boundary of polytope.

The Dehn–Sommerville equations, Upper Bound and Lower Bound theorems were proved in different times for arbitrary triangulations of spheres. However, the generalisation of Theorem 1.3 to arbitrary simplicial spheres remains the main combinatorial conjecture concerning the number of faces.

**Problem 1.14** (*g-conjecture*). Is it true that Theorem 1.3 holds for arbitrary triangulation of  $(n-1)$ -sphere?

## 2. STANLEY–REISNER RINGS: COMBINATORICS AND HOMOLOGICAL ALGEBRA.

The notion of Stanley–Reisner face ring  $\mathbf{k}[K]$  of simplicial complex  $K$  is central in the algebraic study of triangulations. It allows to translate the combinatorics into commutative homological algebra. We review its main properties, emphasising functoriality. Then we introduce the bigraded Tor-algebra  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k})$  through a finite free resolution of  $\mathbb{Z}[K]$  as a module over the polynomial ring. The corresponding bigraded Betti numbers are important combinatorial invariants of  $K$ .

**2.1. Definition of  $\mathbf{k}[K]$ .** Let  $\mathbf{k}[v_1, \dots, v_m]$  be the graded polynomial algebra over  $\mathbf{k} = \mathbb{Z}$  or a field,  $\deg v_i = 2$ . For arbitrary subset  $\sigma = \{i_1, \dots, i_k\} \subset [m]$  denote by  $v_\sigma$  the square-free monomial  $v_{i_1} \dots v_{i_k}$ .

The *face ring* (or the *Stanley–Reisner ring*) of a simplicial complex  $K$  on the vertex set  $[m]$  is the quotient ring

$$\mathbf{k}[K] = \mathbf{k}[v_1, \dots, v_m] / \mathcal{I}_K,$$

where  $\mathcal{I}_K$  is the homogeneous ideal generated by all monomials  $v_\sigma$  such that  $\sigma$  is not a simplex of  $K$ . The ideal  $\mathcal{I}_K$  is called the *Stanley–Reisner ideal* of  $K$ .

**Example 2.1.** 1. Let  $K$  be a 2-dimensional simplicial complex shown on Figure 1. Then

$$\mathcal{I}_K = (v_1 v_5, v_3 v_4, v_1 v_2 v_3, v_2 v_4 v_5).$$




FIGURE 1.

**Exercise 2.2.** Show that every square-free monomial ideal in the polynomial ring has the form  $\mathcal{I}_K$  for some simplicial complex  $K$ .

A *missing face* of  $K$  is a subset  $\sigma \subset [m]$  such that  $\sigma \notin K$ , but every proper subset of  $\sigma$  is a simplex of  $K$ . The Stanley–Reisner ideal  $\mathcal{I}_K$  has basis of monomials  $v_\sigma$  corresponding to missing faces  $\sigma$  of  $K$ .  $K$  is called a *flag complex* if any set of vertices which are pairwise connected spans a simplex of  $K$ . Alternatively,  $K$  is a flag complex if and only if every its missing face has two vertices. The Stanley–Reisner ring  $\mathbf{k}[K]$  is a *quadratic algebra* (i.e. the ideal  $\mathcal{I}_K$  is generated by quadratic monomials) if and only if  $K$  is a flag complex.

Let  $K_1$  and  $K_2$  be two simplicial complexes on the vertex sets  $[m_1]$  and  $[m_2]$  respectively. A set map  $\varphi: [m_1] \rightarrow [m_2]$  is called a *simplicial map* between  $K_1$  and  $K_2$  if  $\varphi(\sigma) \in K_2$  for any  $\sigma \in K_1$ .

**Proposition 2.3.** Let  $\varphi: K_1 \rightarrow K_2$  be a simplicial map. Define the map  $\varphi^*: \mathbf{k}[w_1, \dots, w_{m_2}] \rightarrow \mathbf{k}[v_1, \dots, v_{m_1}]$  by

$$\varphi^*(w_j) := \sum_{i \in \varphi^{-1}(j)} v_i.$$

Then  $\varphi^*$  descends to a homomorphism  $\mathbf{k}[K_2] \rightarrow \mathbf{k}[K_1]$  (which we will also denote by  $\varphi^*$ ).

*Proof.* We have to check that  $\varphi^*(\mathcal{I}_{K_2}) \subset \mathcal{I}_{K_1}$ . Suppose  $\tau = \{j_1, \dots, j_s\} \subset [m_2]$  is not a simplex of  $K_2$ . Then

$$(2.4) \quad \varphi^*(w_{j_1} \cdots w_{j_s}) = \sum_{i_1 \in \varphi^{-1}(j_1), \dots, i_s \in \varphi^{-1}(j_s)} v_{i_1} \cdots v_{i_s}.$$

We claim that  $\sigma = \{i_1, \dots, i_s\}$  is not a simplex of  $K_1$  for any monomial  $v_{i_1} \cdots v_{i_s}$  in the right hand side of the above identity. Indeed, otherwise we would have  $\varphi(\sigma) = \tau \in K_2$  by the definition of simplicial map, which is impossible. Hence, the right hand side of (2.4) is in  $\mathcal{I}_{K_1}$ .  $\square$

Assuming that  $\mathbf{k}$  is a field, for any graded  $\mathbf{k}$ -module  $M = M^0 \oplus M^1 \oplus \dots$  define its *Poincaré series* by

$$F(M; t) = \sum_{i=0}^{\infty} (\dim_{\mathbf{k}} M^i) t^i.$$

**Lemma 2.5** (Stanley). The Poincaré series of  $\mathbf{k}[K^{n-1}]$  can be calculated as

$$F(\mathbf{k}[K^{n-1}]; t) = \sum_{i=-1}^{n-1} \frac{f_i t^{2(i+1)}}{(1-t^2)^{i+1}} = \frac{h_0 + h_1 t^2 + \dots + h_n t^{2n}}{(1-t^2)^n},$$

where  $(f_0, \dots, f_{n-1})$  is the  $f$ -vector and  $(h_0, \dots, h_n)$  is the  $h$ -vector of  $K$ .

*Proof.* A monomial in  $\mathbf{k}[K]$  has the form  $v_{i_1}^{\alpha_1} \cdots v_{i_{k+1}}^{\alpha_{k+1}}$  where  $\{i_1, \dots, i_{k+1}\}$  is a simplex of  $K^{n-1}$  and  $\alpha_1, \dots, \alpha_{k+1}$  are some positive integers. Thus, every  $k$ -simplex of  $K$  contributes the summand  $\frac{t^{2(k+1)}}{(1-t^2)^{k+1}}$  to the Poincaré series, which proves the first identity. The second identity is an obvious corollary of (1.1).  $\square$

**Example 2.6.** 1. Let  $K = \Delta^{n-1}$ . Then  $f_i = \binom{n}{i+1}$  for  $-1 \leq i \leq n-1$ ,  $h_0 = 1$  and  $h_i = 0$  for  $i > 0$ . We have  $\mathbf{k}[\Delta^{n-1}] = \mathbf{k}[v_1, \dots, v_n]$  and  $F(\mathbf{k}[\Delta^{n-1}]; t) = (1 - t^2)^{-n}$ , which agrees with Lemma 2.5.

2. Let  $K$  be the boundary of an  $n$ -simplex. Then  $h_i = 1$ ,  $i = 0, 1, \dots, n$ , and  $\mathbf{k}[K] = \mathbf{k}[v_1, \dots, v_{n+1}]/(v_1 v_2 \cdots v_{n+1})$ . By Lemma 2.5,

$$F(\mathbf{k}[K]; t) = \frac{1 + t^2 + \cdots + t^{2n}}{(1 - t^2)^n}.$$

**2.2. Cohen–Macaulay algebras.** Suppose now that  $A = \bigoplus_{i \geq 0} A^i$  is a finitely-generated commutative graded algebra over  $\mathbf{k}$ . We assume that  $A$  is connected ( $A^0 = \mathbf{k}$ ) and has only even-degree graded components, so it is commutative as either ordinary or graded algebra. We denote by  $A_+$  the positive-degree part of  $A$  and by  $\mathcal{H}(A_+)$  the set of homogeneous elements in  $A_+$ . The *Krull dimension* of  $A$ , denoted  $\dim A$ , is the maximal number of algebraically independent elements of  $A$ .

A sequence  $t_1, \dots, t_n$  of algebraically independent homogeneous elements of  $A$  is called an *hsop* (homogeneous system of parameters) if  $A$  is a finitely-generated  $\mathbf{k}[t_1, \dots, t_n]$ -module (equivalently,  $A/(t_1, \dots, t_n)$  has finite dimension as a  $\mathbf{k}$ -vector space).

**Lemma 2.7** (Nöther normalisation lemma). *Any finitely-generated graded algebra  $A$  over a field  $\mathbf{k}$  admits an hsop. If  $\mathbf{k}$  is of zero characteristic and  $A$  is generated by degree-two elements, then a degree-two hsop can be chosen.*

A degree-two hsop is called an *lsop* (linear system of parameters).

A sequence  $\mathbf{t} = t_1, \dots, t_k$  of elements of  $\mathcal{H}(A_+)$  is called a *regular sequence* if  $t_{i+1}$  is not a zero divisor in  $A/(t_1, \dots, t_i)$  for  $0 \leq i < k$ .

**Exercise 2.8.** Prove that any regular sequence consists of algebraically independent elements, so it generates a polynomial subring in  $A$ . Show further that  $\mathbf{t}$  is a regular sequence if and only if  $A$  is a free  $\mathbf{k}[t_1, \dots, t_k]$ -module.

Algebra  $A$  is called *Cohen–Macaulay* if it admits a regular hsop  $\mathbf{t}$ . It follows that  $A$  is Cohen–Macaulay if and only if it is free finitely generated module over its polynomial subring. If  $\mathbf{k}$  is a field of zero characteristic and  $A$  is generated by degree-two elements, then one can choose  $\mathbf{t}$  to be an lsop. In this case the following formula for the Poincaré series of  $A$  holds

$$F(A; t) = \frac{F(A/(t_1, \dots, t_n); t)}{(1 - t^2)^n},$$

where  $F(A/(t_1, \dots, t_n); t)$  is a polynomial.

A simplicial complex  $K$  is called *Cohen–Macaulay* (over  $\mathbf{k}$ ) if its face ring  $\mathbf{k}[K]$  is Cohen–Macaulay.

**Exercise 2.9.** Prove that the Krull dimension of  $\mathbf{k}[K^{n-1}]$  equals  $n$ . Compare the above formula for  $F(\mathbf{k}[K], \mathbf{k})$  with that of Lemma 2.5.

**Example 2.10.** Let  $K = \partial\Delta^2$  be the boundary of a 2-simplex. Then  $\mathbf{k}[K] = \mathbf{k}[v_1, v_2, v_3]/(v_1 v_2 v_3)$ . The elements  $v_1, v_2 \in \mathbf{k}[K]$  are algebraically independent, but do not form an hsop, since  $\mathbf{k}[K]/(v_1, v_2) \cong \mathbf{k}[v_3]$  is not finite-dimensional as

a  $\mathbf{k}$ -space. On the other hand, the elements  $t_1 = v_1 - v_3$ ,  $t_2 = v_2 - v_3$  of  $\mathbf{k}[K]$  form an hso, since  $\mathbf{k}[K]/(t_1, t_2) \cong \mathbf{k}[t]/t^3$ . It is easy to see that  $\mathbf{k}[K]$  is a free  $\mathbf{k}[t_1, t_2]$ -module with one 0-dimensional generator 1, one 1-dimensional generator  $v_1$ , and one 2-dimensional generator  $v_1^2$ . Thus,  $\mathbf{k}[K]$  is Cohen–Macaulay and  $(t_1, t_2)$  is a regular sequence.

**Proposition 2.11** (Stanley). *If  $K^{n-1}$  is a Cohen–Macaulay simplicial complex, then  $\mathbf{h}(K^{n-1}) = (h_0, \dots, h_n)$  is an  $M$ -vector.*

*Proof.* Let  $t_1, \dots, t_n$  be a regular sequence of degree-two elements of  $\mathbf{k}[K]$ . Then  $A = \mathbf{k}[K]/(t_1, \dots, t_n)$  is a graded algebra generated by degree-two elements, and  $\dim_{\mathbf{k}} A^{2i} = h_i$ . Now the result follows from Theorem 1.7.  $\square$

For arbitrary simplex  $\sigma \in K$  define its *link* and *star* as subcomplexes

$$\begin{aligned} \text{link}_K \sigma &= \{\tau \in K : \sigma \cup \tau \in K, \sigma \cap \tau = \emptyset\}; \\ \text{star}_K \sigma &= \{\tau \in K : \sigma \cup \tau \in K\}. \end{aligned}$$

For any vertex  $v \in K$  the subcomplex  $\text{star}_K v$  can be identified with the cone over  $\text{link}_K v$ . The polyhedron  $|\text{star}_K v|$  consists of all faces of  $|K|$  that contain  $v$ . We omit the subscript  $K$  whenever the context allows.

The following fundamental theorem characterises Cohen–Macaulay complexes combinatorially.

**Theorem 2.12** (Reisner). *A simplicial complex  $K$  is Cohen–Macaulay over  $\mathbf{k}$  if and only if for any simplex  $\sigma \in K$  (including  $\sigma = \emptyset$ ) and  $i < \dim(\text{link } \sigma)$  it holds that  $\tilde{H}_i(\text{link } \sigma; \mathbf{k}) = 0$ .*

The standard  $PL$  topology techniques allow to reformulate the above theorem in purely topological terms.

**Proposition 2.13** (Munkres).  *$K^{n-1}$  is Cohen–Macaulay over  $\mathbf{k}$  if and only if for arbitrary point  $x \in |K|$  it holds that  $\tilde{H}_i(|K|; \mathbf{k}) = H_i(|K|, |K| \setminus x; \mathbf{k}) = 0$  for  $i < n - 1$ .*

**Corollary 2.14.** *A triangulation of sphere is a Cohen–Macaulay complex.*

Now by Theorem 2.11 one concludes that the  $h$ -vector of a simplicial sphere is an  $M$ -vector. This argument was used by Stanley to extend the Upper Bound Theorem (Theorem 1.10) to simplicial spheres.

**Corollary 2.15** (UBT for spheres, Stanley). *For arbitrary simplicial  $(n - 1)$ -sphere  $K^{n-1}$  with  $m$  vertices it holds that*

$$f_i(K^{n-1}) \leq f_i(C^n(m)) \quad \text{for } i = 1, \dots, n - 1.$$

*Proof.* Due to Exercise 1.12, the UBT is equivalent to the inequalities

$$h_i(K^{n-1}) \leq \binom{m-n+i-1}{i}, \quad 0 \leq i < \left\lfloor \frac{n}{2} \right\rfloor.$$

Since  $\mathbf{h}(K^{n-1})$  is an  $M$ -vector, there exists a graded algebra  $A = A^0 \oplus A^2 \oplus \dots \oplus A^{2n}$  generated by degree-two elements such that  $\dim_{\mathbf{k}} A^{2i} = h_i$  (Theorem 1.7). In particular,  $\dim_{\mathbf{k}} A^2 = h_1 = m - n$ . Since  $A$  is generated by  $A^2$ , the number  $h_i$  cannot exceed the total number of monomials of degree  $i$  in  $(m - n)$  variables. The latter is exactly  $\binom{m-n+i-1}{i}$ .  $\square$

**2.3. Resolutions and Tor-algebras.** Let  $M$  be a finitely-generated graded  $\mathbf{k}[v_1, \dots, v_m]$ -module. A *free resolution* of  $M$  is an exact sequence (2.16)

$$\dots \xrightarrow{d} R^{-i} \xrightarrow{d} \dots \xrightarrow{d} R^{-1} \xrightarrow{d} R^0 \longrightarrow M \longrightarrow 0,$$

where the  $R^{-i}$  are finitely-generated free modules and the maps  $d$  are degree-preserving. The minimal number  $h$  for which there exists a free resolution (2.16) with  $R^{-i} = 0$  for  $i > h$  is called the *homological dimension* of  $M$  and denoted  $\text{hd } M$ . By the Hilbert syzygy theorem,  $\text{hd } M \leq m$ . A resolution (2.16) can be written as a free *bigraded differential  $\mathbf{k}$ -module*  $[R, d]$ , where  $R = \bigoplus R^{-i,j}$ ,  $R^{-i,j} := (R^{-i})^j$  and  $d: R^{-i,j} \rightarrow R^{-i+1,j}$ . The bigraded cohomology module  $H[R, d]$  has  $H^{-i,k}[R, d] = 0$  for  $i > 0$  and  $H^{0,k}[R, d] = M^k$ . Let  $[M, 0]$  be the bigraded module with  $M^{-i,k} = 0$  for  $i > 0$ ,  $M^{0,k} = M^k$ , and zero differential. Then the resolution (2.16) determined a map  $[R, d] \rightarrow [M, 0]$  inducing an isomorphism in cohomology.

**Exercise 2.17.** Show that the Poincaré series of  $M$  can be calculated from any free resolution (2.16) as follows. Suppose that  $R^{-i}$  has rank  $q_i$  with free generators in degrees  $d_{1i}, \dots, d_{q_i i}$ ,  $i = 1, \dots, h$ . Then

$$(2.18) \quad F(M; t) = (1 - t^2)^{-m} \sum_{i=0}^h (-1)^i (t^{d_{1i}} + \dots + t^{d_{q_i i}}).$$

**Example 2.19** (Koszul resolution). Let  $M = \mathbf{k}$ . The  $\mathbf{k}[v_1, \dots, v_m]$ -module structure on  $\mathbf{k}$  is defined via the map  $\mathbf{k}[v_1, \dots, v_m] \rightarrow \mathbf{k}$  that sends each  $v_i$  to 0. Let  $\Lambda[u_1, \dots, u_m]$  denote the exterior algebra on  $m$  generators. Turn the tensor product  $R = \Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[v_1, \dots, v_m]$  (here and below we use  $\otimes$  for  $\otimes_{\mathbf{k}}$ ) into a *differential bigraded algebra* by setting

$$(2.20) \quad \begin{aligned} \text{bideg } u_i &= (-1, 2), & \text{bideg } v_i &= (0, 2), \\ du_i &= v_i, & dv_i &= 0, \end{aligned}$$

and requiring that  $d$  be a derivation of algebras. An explicit construction of cochain homotopy shows that  $H^{-i}[R, d] = 0$  for  $i > 0$  and  $H^0[R, d] = \mathbf{k}$ . Since  $\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[v_1, \dots, v_m]$  is a free  $\mathbf{k}[v_1, \dots, v_m]$ -module, it determines a free resolution of  $\mathbf{k}$ . This resolution is known as the *Koszul resolution*. Its expanded form is as follows:

$$\begin{aligned} 0 \rightarrow \Lambda^m[u_1, \dots, u_m] \otimes \mathbf{k}[v_1, \dots, v_m] &\longrightarrow \dots \\ &\longrightarrow \Lambda^1[u_1, \dots, u_m] \otimes \mathbf{k}[v_1, \dots, v_m] \longrightarrow \mathbf{k}[v_1, \dots, v_m] \longrightarrow \mathbf{k} \rightarrow 0, \end{aligned}$$

where  $\Lambda^i[u_1, \dots, u_m]$  is the submodule of  $\Lambda[u_1, \dots, u_m]$  spanned by monomials of length  $i$ .

Let  $N$  be another module; then applying the functor  $\otimes_{\mathbf{k}[v_1, \dots, v_m]} N$  to resolution  $[R, d]$  we get a homomorphism of differential modules

$$[R \otimes_{\mathbf{k}[v_1, \dots, v_m]} N, d] \rightarrow [M \otimes_{\mathbf{k}[v_1, \dots, v_m]} N, 0],$$



which in general does not induce an isomorphism in cohomology. The  $(-i)$ th cohomology module of the cochain complex

$$\dots \longrightarrow R^{-i} \otimes_{\mathbf{k}[v_1, \dots, v_m]} N \longrightarrow \dots \longrightarrow R^0 \otimes_{\mathbf{k}[v_1, \dots, v_m]} N \longrightarrow 0$$

is denoted  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i}(M, N)$ . Thus,

$$\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i}(M, N) := \frac{\mathrm{Ker}[d: R^{-i} \otimes_{\mathbf{k}[v_1, \dots, v_m]} N \rightarrow R^{-i+1} \otimes_{\mathbf{k}[v_1, \dots, v_m]} N]}{d(R^{-i-1} \otimes_{\mathbf{k}[v_1, \dots, v_m]} N)}.$$

Since all the  $R^{-i}$  and  $N$  are graded modules, we actually have the *bigraded*  $\mathbf{k}$ -module

$$\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(M, N) = \bigoplus_{i,j} \mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i,j}(M, N).$$

The following properties of  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i}(M, N)$  are well known.

**Proposition 2.21.** (a) *The module  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i}(M, N)$  does not depend, up to isomorphism, on a choice of resolution (2.16).*

(b) *Both  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i}(\cdot, N)$  and  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i}(M, \cdot)$  are covariant functors.*

(c)  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^0(M, N) \cong M \otimes_{\mathbf{k}[v_1, \dots, v_m]} N$ .

(d)  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i}(M, N) \cong \mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i}(N, M)$ .

Now put  $M = \mathbf{k}[K]$  and  $N = \mathbf{k}$ . Since  $\deg v_i = 2$ , we have

$$\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k}) = \bigoplus_{i,j=0}^m \mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i,2j}(\mathbf{k}[K], \mathbf{k})$$

(i.e.  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k})$  is non-zero only in even second degrees). Define the *bigraded Betti numbers* of  $\mathbf{k}[K]$  by

$$(2.22) \quad \beta^{-i,2j}(\mathbf{k}[K]) := \dim_{\mathbf{k}} \mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i,2j}(\mathbf{k}[K], \mathbf{k}), \quad 0 \leq i, j \leq m.$$

We also set

$$\beta^{-i}(\mathbf{k}[K]) = \dim_{\mathbf{k}} \mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i}(\mathbf{k}[K], \mathbf{k}) = \sum_j \beta^{-i,2j}(\mathbf{k}[K]).$$

**Example 2.23.** Let  $K$  be the boundary of a square. Then

$$\mathbf{k}[K] \cong \mathbf{k}[v_1, \dots, v_4]/(v_1v_3, v_2v_4).$$

Let us construct a resolution of  $\mathbf{k}[K]$  and calculate the corresponding bigraded Betti numbers. The module  $R^0$  has one generator 1 (of degree 0), and the map  $R^0 \rightarrow \mathbf{k}[K]$  is the quotient projection. Its kernel is the ideal  $\mathcal{I}_{K^1}$ , generated by two monomials  $v_1v_3$  and  $v_2v_4$ . Take  $R^{-1}$  to be the free module on two 4-dimensional generators, denoted  $v_{13}$  and  $v_{24}$ , and define  $d: R^{-1} \rightarrow R^0$  by sending  $v_{13}$  to  $v_1v_3$  and  $v_{24}$  to  $v_2v_4$ . Its kernel is generated by one element  $v_2v_4v_{13} - v_1v_3v_{24}$ . Hence,  $R^{-2}$  has one generator of degree 8, say  $a$ , and the map  $d: R^{-2} \rightarrow R^{-1}$  is injective and sends  $a$  to  $v_2v_4v_{13} - v_1v_3v_{24}$ . Thus, we have resolution

$$0 \longrightarrow R^{-2} \longrightarrow R^{-1} \longrightarrow R^0 \longrightarrow M \longrightarrow 0,$$

where  $\mathrm{rank} R^0 = \beta^{0,0}(\mathbf{k}[K]) = 1$ ,  $\mathrm{rank} R^{-1} = \beta^{-1,4} = 2$ ,  $\mathrm{rank} R^{-2} = \beta^{-2,8} = 1$ .

The Betti numbers  $\beta^{-i,2j}(\mathbf{k}[K])$  are important combinatorial invariants of simplicial complex  $K$ . The following result expresses them in terms of homology groups of subcomplexes of  $K$ .

**Theorem 2.24** (Hochster). *We have*

$$\beta^{-i,2j}(\mathbf{k}[K]) = \sum_{\omega \subset [m]: |\omega|=j} \dim_{\mathbf{k}} \tilde{H}^{j-i-1}(K_{\omega}; \mathbf{k}),$$

where  $K_{\omega}$  is the full subcomplex of  $K$  spanned by  $\omega$ . We assume  $\tilde{H}^{-1}(\emptyset) = \mathbf{k}$  above.

The original Hochster's proof of this theorem uses rather complicated combinatorial and commutative algebra techniques. Later we give another topological proof.

**Exercise 2.25.** Calculate the bigraded Betti numbers from Example 2.23 using Hochster theorem.

Now let us consider the differential bigraded algebra  $[\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[K], d]$  with  $d$  defined as in (2.20).

**Lemma 2.26.**  $\text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k})$  is an algebra in a canonical way, and there is an isomorphism of algebras

$$\text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k}) \cong H[\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[K], d].$$

*Proof.* Using the Koszul resolution  $[\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[v_1, \dots, v_m], d]$  in the definition of  $\text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}, \mathbf{k}[K])$ , we calculate

$$\begin{aligned} \text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k}) &\cong \text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}, \mathbf{k}[K]) \\ &= H[\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[v_1, \dots, v_m] \otimes_{\mathbf{k}[v_1, \dots, v_m]} \mathbf{k}[K]] \\ &\cong H[\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[K]]. \end{aligned}$$

□

The bigraded algebra  $\text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k})$  is called the *Tor-algebra* of simplicial complex  $K$ .

**Lemma 2.27.** A simplicial map  $\varphi: K_1 \rightarrow K_2$  between two simplicial complexes on the vertex sets  $[m_1]$  and  $[m_2]$  respectively induces a homomorphism

$$(2.28) \quad \varphi_t^*: \text{Tor}_{\mathbf{k}[w_1, \dots, w_{m_2}]}(\mathbf{k}(K_2), \mathbf{k}) \rightarrow \text{Tor}_{\mathbf{k}[v_1, \dots, v_{m_1}]}(\mathbf{k}(K_1), \mathbf{k})$$

of the corresponding Tor-algebras.

*Proof.* This follows directly from Propositions 2.3 and 2.21 (b). □

**2.4. Gorenstein\* complexes.** Cohen–Macaulay complexes may be characterised by means of Betti numbers as follows.

**Exercise 2.29.**  $K^{n-1}$  is a Cohen–Macaulay complex if and only if  $\beta^{-i}(\mathbf{k}[K]) = 0$  for  $i > m - n$ . In this case  $\beta^{-(m-n)}(\mathbf{k}[K]) \neq 0$ .

A Cohen–Macaulay complex  $K^{n-1}$  on the set  $[m]$  is called *Gorenstein* if  $\beta^{-(m-n)}(\mathbf{k}[K]) = 1$ , that is,  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-(m-n)}(\mathbf{k}[K], \mathbf{k}) \cong \mathbf{k}$ . If additionally  $K$  is not a cone over another simplicial complex then it is called *Gorenstein\**. In the latter case we have

$$\beta^{-(m-n)}(\mathbf{k}[K]) = \beta^{-(m-n), 2m}(\mathbf{k}[K]) = 1.$$

The following result is similar to Reisner’s theorem 2.12 and characterises Gorenstein\* simplicial complexes.

**Theorem 2.30** (Stanley). *A simplicial complex  $K$  is Gorenstein\* over  $\mathbf{k}$  if and only if for any simplex  $\sigma \in K$  (including  $\sigma = \emptyset$ ) the subcomplex  $\mathrm{link} \sigma$  has the homology of a sphere of dimension  $\dim(\mathrm{link} \sigma)$ .*

In particular, simplicial spheres and simplicial homology spheres (triangulated manifolds with the homology of a sphere) are Gorenstein\* complexes. However, the Gorenstein\* property does not guarantee a complex to be a triangulated manifold. Gorenstein complexes are known to topologists as “generalised homology spheres”.

A graded commutative finite-dimensional connected  $\mathbf{k}$ -algebra  $H = \bigoplus_{i=0}^d H^i$  is called a *Poincaré algebra* if  $\mathbf{k}$ -linear maps  $H^i \rightarrow \mathrm{Hom}_{\mathbf{k}}(H^{d-i}, H^d)$ ,  $a \mapsto \varphi_a$ ,  $\varphi_a(b) = ab$  are isomorphisms for all  $i = 0, \dots, d$ . The following is a corollary of Avramov–Golod theorem [1, Thm. 3.4.5].

**Theorem 2.31.** *A simplicial complex is Gorenstein if and only if its Tor-algebra is Poincaré.*

It is easy to see that the Poincaré algebra structure respects the bigrading, whence the next result follows.

**Corollary 2.32.** *Suppose  $K^{n-1}$  is a Gorenstein\* complex on  $[m]$ . Then the following identities hold for the Poincaré series of Tor-algebra and for the bigraded Betti numbers:*

$$\begin{aligned} F(\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-i}(\mathbf{k}[K], \mathbf{k}); t) &= t^{2m} F(\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}^{-(m-n)+i}(\mathbf{k}[K], \mathbf{k}); \tfrac{1}{t}); \\ \beta^{-i, 2j}(\mathbf{k}[K]) &= \beta^{-(m-n)+i, 2(m-j)}(\mathbf{k}[K]), \quad i = 0, \dots, m-n, \quad j = 0, \dots, m. \end{aligned}$$

**Exercise 2.33.** Deduce from the above that if  $K^{n-1}$  is Gorenstein\* then

$$F(\mathbf{k}[K], t) = (-1)^n F(\mathbf{k}[K], \tfrac{1}{t}).$$

Deduce further that the Dehn–Sommerville relations  $h_i = h_{n-i}$ ,  $0 \leq i \leq n$ , hold for arbitrary Gorenstein\* complex  $K^{n-1}$ .

We see that the class of Gorenstein\* complexes is in a sense “the best possible algebraic approximation” to triangulated spheres. As it was conjectured by Stanley, the  $g$ -theorem may continue to hold for Gorenstein\* complexes.

Later we deduce the generalised Dehn–Sommerville equations for triangulated manifolds as a consequence of the bigraded Poincaré duality for moment-angle complexes. In particular, this gives the following short form of the equations in terms of the  $h$ -vector:

$$h_{n-i} - h_i = (-1)^i (\chi(K^{n-1}) - \chi(S^{n-1})) \binom{n}{i}, \quad i = 0, 1, \dots, n.$$

Here  $\chi(K^{n-1}) = f_0 - f_1 + \dots + (-1)^{n-1}f_{n-1} = 1 + (-1)^{n-1}h_n$  is the Euler characteristic of  $K^{n-1}$  and  $\chi(S^{n-1}) = 1 + (-1)^{n-1}$  is that of a sphere. Note that the above equations reduce to the classical  $h_{n-i} = h_i$  in the case when  $K$  is a simplicial sphere or has odd dimension. Note also that a triangulated manifold is not Gorenstein\*, or even Cohen–Macaulay, in general.

### 3. DAVIS–JANUSZKIEWICZ SPACES AND MOMENT-ANGLE MANIFOLDS AND COMPLEXES.

Here we study different "topological models" for the algebraic objects introduced before. The Davis–Januszkiewicz space  $DJ(K)$  has the cohomology isomorphic to the Stanley–Reisner ring  $\mathbf{k}[K]$ . A Borel construction type model of such space is due to Davis and Januszkiewicz [3], whence the name comes. It also appeared in different disguises in the work of Hattori–Masuda on torus manifolds. In our approach we use another model for  $DJ(K)$ , defined through a simple colimit (or "nested union") of nice building blocks. It is homotopy equivalent to the Davis–Januszkiewicz model. The building block space depends on the coefficients  $\mathbf{k}$  (e.g., it is  $\mathbb{C}P^\infty$  if  $\mathbf{k} = \mathbb{Z}$ , and  $\mathbb{R}P^\infty$  if  $\mathbf{k} = \mathbb{Z}/2$ ). The term "moment-angle complex" refers to a special bigraded cellular decomposition of the universal space  $\mathcal{Z}_K$  acted on by the torus. This space was also introduced by Davis and Januszkiewicz. It was defined as the fibre of a bundle with total space  $DJ(K)$  and base  $BT^m$ , but it has also many other interesting interpretations. The moment-angle complex  $\mathcal{Z}_K$  is a manifold provided that  $K$  is a triangulation of a sphere. At the end we discuss how the spaces  $DJ(K)$  and  $\mathcal{Z}_K$  are related to toric varieties and fans, Davis–Januszkiewicz (quasi)toric manifolds and characteristic functions, and Hattori–Masuda torus manifolds and multifans.

**3.1. Definitions and main properties.** Here we assume  $\mathbf{k} = \mathbb{Z}$ , unless otherwise specified.

The classifying space for the circle  $S^1$  can be identified with the infinite-dimensional projective space  $\mathbb{C}P^\infty$ . It has a canonical cell decomposition with one cell  $B^{2k}$  in each even dimension. The classifying space  $BT^m$  of the  $m$ -torus is thus the product of  $m$  copies of  $\mathbb{C}P^\infty$ . The cohomology of  $BT^m$  is the polynomial ring  $\mathbb{Z}[v_1, \dots, v_m]$ ,  $\deg v_i = 2$ . The total space  $ET^m$  of the universal principal  $T^m$ -bundle over  $BT^m$  can be identified with the product of  $m$  infinite-dimensional spheres.

For arbitrary subset  $\omega \subset [m]$  define the subproduct

$$BT^\omega := \{(x_1, \dots, x_m) \in BT^m : x_i = * \text{ if } i \notin \omega\}.$$

For arbitrary simplicial complex  $K$  on  $[m]$  define the *Davis–Januszkiewicz space* as the following cellular subcomplex:

$$DJ(K) := \bigcup_{\sigma \in K} BT^\sigma \subset BT^m.$$

**Proposition 3.1.** *The cohomology of  $DJ(K)$  is isomorphic to the Stanley–Reisner ring  $\mathbb{Z}[K]$ . Moreover, the inclusion of cellular complexes  $i: DJ(K) \hookrightarrow$*

$BT^m$  induces the quotient epimorphism

$$i^*: \mathbb{Z}[v_1, \dots, v_m] \rightarrow \mathbb{Z}[K] = \mathbb{Z}[v_1, \dots, v_m]/\mathcal{I}_K$$

in the cohomology.

*Proof.* A monomial  $v_{i_1}^{k_1} \dots v_{i_p}^{k_p}$  represents the cellular cochain  $(B_{i_1}^{2k_1} \dots B_{i_p}^{2k_p})^*$  in  $C^*(BT^m)$ . Under the cochain homomorphism induced by the inclusion  $DJ(K) \subset BT^m$  it maps identically if  $\{i_1, \dots, i_p\} \in K$  and to zero otherwise, whence the statement follows.  $\square$

It is convenient to realise the torus  $T^m$  as a subspace in  $\mathbb{C}^m$ :

$$T^m = \{(z_1, \dots, z_m) \in \mathbb{C}^m : |z_i| = 1, \quad i = 1, \dots, m\}.$$

It is contained in the *unit polydisc*

$$(D^2)^m := \{(z_1, \dots, z_m) \in \mathbb{C}^m : |z_i| \leq 1, \quad i = 1, \dots, m\}.$$

For arbitrary subset  $\omega \subset [m]$  define

$$B_\omega := \{(z_1, \dots, z_m) \in (D^2)^m : |z_i| = 1 \text{ if } i \notin \omega\}.$$

Obviously,  $B_\omega$  is homeomorphic to  $(D^2)^{|\omega|} \times T^{m-|\omega|}$ . Given a simplicial complex  $K$  on  $[m]$ , define the *moment-angle complex*  $\mathcal{Z}_K$  by

$$(3.2) \quad \mathcal{Z}_K := \bigcup_{\sigma \in K} B_\sigma \subset (D^2)^m.$$

*Remark.* The above constructions of  $DJ(K)$  and  $\mathcal{Z}_K$  are examples of *colimits* of diagrams of topological spaces over the *face category* of  $K$  (objects are simplices and morphisms are inclusions). In the first case the diagram assigns the space  $BT^\sigma$  to a simplex  $\sigma$ ; its colimit is  $DJ(K)$ . The second diagram assigns  $B_\sigma$  to  $\sigma$ ; its colimit is  $\mathcal{Z}_K$ .

The torus  $T^m$  acts on  $(D^2)^m$  coordinatewise, and each subspace  $B_\omega$  is invariant under this action. Therefore, the space  $\mathcal{Z}_K$  itself is acted on by the torus. The quotient  $(D^2)^m/T^m$  can be identified with the *unit  $m$ -cube*:

$$I^m := \{(y_1, \dots, y_m) \in \mathbb{R}^m : 0 \leq y_i \leq 1, \quad i = 1, \dots, m\}.$$

The quotient  $B_\omega/T^m$  is then the following  $|\omega|$ -face of  $I^m$ :

$$C_\omega := \{(y_1, \dots, y_m) \in I^m : y_i = 1 \text{ if } i \notin \omega\}.$$

Thus, the whole quotient  $\mathcal{Z}_K/T^m$  is identified with a certain cubical subcomplex in  $I^m$ , which we denote  $\text{cc}(K)$ .

**Lemma 3.3.** *The cubical complex  $\text{cc}(K)$  is PL-homeomorphic to  $|\text{cone } K|$ .*

*Proof.* Let  $K'$  denote the barycentric subdivision of  $K$  (the vertices of  $K'$  correspond to non-empty simplices  $\sigma$  of  $K$ ). We define a *PL* embedding  $i_c: \text{cone } K' \hookrightarrow I^m$  by sending each vertex  $\sigma$  to the vertex  $(\varepsilon_1, \dots, \varepsilon_m) \in I^m$ , where  $\varepsilon_i = 0$  if  $i \in \sigma$  and  $\varepsilon_i = 1$  otherwise, sending the cone vertex to  $(1, \dots, 1) \in I^m$ , and extending linearly on the simplices of  $\text{cone } K'$ . The barycentric subdivision of a face  $\sigma \in K$  is a subcomplex in  $K'$ , which we denote  $K'|_\sigma$ . Under the map  $i_c$  the subcomplex  $\text{cone } K'|_\sigma$  maps onto the face  $C_\sigma \subset I^m$ .

Thus, the whole complex  $\text{cone } K'$  maps homeomorphically onto  $\text{cc}(K)$ , whence the proof follows.  $\square$

**Example 3.4.** If  $K = \Delta^{m-1}$  is the whole simplex on  $[m]$ , then  $\text{cc}(K)$  is the whole cube  $I^m$ , and the above constructed  $PL$ -homeomorphism between  $\text{cone}(\Delta^{m-1})'$  to  $I^m$  defines the *canonical triangulation* of  $I^m$ .

**Lemma 3.5.** *Suppose that  $K$  is a triangulation of a sphere:  $|K| \cong S^{n-1}$ . Then  $\mathcal{Z}_K$  is an  $(m+n)$ -dimensional manifold.*

*Proof.* The space  $|\text{cone } K'|$  has a canonical *face structure* whose facets (codimension-one faces) are  $F_i := \text{star}_{K'}\{i\}$ ,  $i = 1, \dots, m$ , and  $i$ -faces are non-empty intersections of  $i$ -tuples of facets. Since  $K$  is a triangulation of a sphere,  $|\text{cone } K'|$  is an  $n$ -ball. Every point in  $|\text{cone } K'|$  has a neighbourhood homeomorphic to an open subset in  $I^n$  (or  $\mathbb{R}_+^n$ ) with the homeomorphism preserving the dimension of faces. By the definition, this displays  $|\text{cone } K'|$  as a *manifold with corners*. Having identified  $|\text{cone } K'|$  with  $\text{cc}(K)$  and further  $\text{cc}(K)$  with  $\mathcal{Z}_K/T^m$ , we see that every point in  $\mathcal{Z}_K$  lies in a neighbourhood homeomorphic to an open subset in  $(D^2)^n \times T^{m-n}$  and thus in  $\mathbb{R}^{m+n}$ .  $\square$

Suppose  $P^n$  is a *simple*  $n$ -polytope, that is, every vertex of  $P$  is contained in exactly  $n$  facets. Define  $K_P$  to be the boundary complex of the dual simplicial polytope. Then the face structure in  $\text{cone } K'_P$  is the same as that of  $P^n$ .

**Exercise 3.6.** Show that the isotropy subgroup of a point  $x \in \text{cc}(K)$  under the  $T^m$ -action on  $\mathcal{Z}_K$  is the coordinate subtorus

$$T(x) := \{(z_1, \dots, z_m) \in T^m : z_i = 1 \text{ if } x \notin F_i\}.$$

In particular, the action is free over the interior (that is, around the cone point) of  $\text{cc}(K) \cong |\text{cone } K'|$ .

It follows that the moment-angle complex can be identified with the quotient

$$\mathcal{Z}_K = (T^m \times |\text{cone } K'|)/\sim,$$

where  $(t_1, x) \sim (t_2, y)$  if and only if  $x = y$  and  $t_1 t_2^{-1} \in T(x)$ . In the case  $K = K_P$  we may write  $(T^m \times P^n)/\sim$  instead. The latter  $T^m$ -manifold is the one introduced by Davis and Januszkiewicz [3], which thereby coincides with our moment-angle complex.

For arbitrary  $T^m$ -space  $X$  define the *Borel construction* (also known as the *homotopy quotient* or *associated bundle*) as the identification space

$$ET^m \times_{T^m} X := ET^m \times X / \sim,$$

where  $(e, x) \sim (eg, g^{-1}x)$  for any  $e \in ET^m$ ,  $x \in X$ ,  $g \in T^m$ .

The projection  $(e, x) \rightarrow e$  identifies  $ET^m \times_{T^m} X$  with the total space of a bundle  $ET^m \times_{T^m} X \rightarrow BT^m$  with fibre  $X$  and structure group  $T^m$ .

In the sequel we denote the Borel construction  $ET^m \times_{T^m} X$  by  $B_T X$ .

In particular, for any simplicial complex  $K$  on  $m$  vertices we have the Borel construction  $B_T \mathcal{Z}_K$  and the bundle  $p: B_T \mathcal{Z}_K \rightarrow BT^m$  with fibre  $\mathcal{Z}_K$ . Davis and Januszkiewicz showed that the cohomology of  $B_T \mathcal{Z}_K$  is isomorphic to the Stanley–Reisner ring  $\mathbb{Z}[K]$ , and  $p^*: \mathbb{Z}[v_1, \dots, v_m] \rightarrow \mathbb{Z}[K]$  is the quotient projection. Thus, the space  $B_T \mathcal{Z}_K$  provides another topological model for the

Stanley–Reisner ring. However, the following result shows that the two models  $DJ(K)$  and  $B_T \mathcal{Z}_K$  are homotopy equivalent.

**Theorem 3.7.** *There is a deformation retraction  $B_T \mathcal{Z}_K \rightarrow DJ(K)$  such that the diagram*

$$\begin{array}{ccc} B_T \mathcal{Z}_K & \xrightarrow{p} & BT^m \\ \downarrow & & \parallel \\ DJ(K) & \xrightarrow{i} & BT^m \end{array}$$

*is commutative.*

*Proof.* We have  $\mathcal{Z}_K = \bigcup_{\sigma \in K} B_\sigma$ , and each  $B_\sigma$  is  $T^m$ -invariant. Hence, there is the corresponding decomposition of the Borel construction:

$$B_T \mathcal{Z}_K = ET^m \times_{T^m} \mathcal{Z}_K = \bigcup_{\sigma \in K} ET^m \times_{T^m} B_\sigma.$$

Suppose  $|\sigma| = s$ . Then  $B_\sigma \cong (D^2)^s \times T^{m-s}$ , so we have  $ET^m \times_{T^m} B_\sigma \cong (ET^s \times_{T^s} (D^2)^s) \times ET^{m-s}$ . The space  $ET^s \times_{T^s} (D^2)^s$  is the total space of a  $(D^2)^s$ -bundle over  $BT^s$ , and  $ET^{m-s}$  is contractible. It follows that there is a deformation retraction  $ET^m \times_{T^m} B_\sigma \rightarrow BT^\sigma$ . These homotopy equivalences corresponding to different simplices fit together to yield a required homotopy equivalence between  $p: B_T \mathcal{Z}_K \rightarrow BT^m$  and  $i: DJ(K) \hookrightarrow BT^m$ .  $\square$

Below we denote by  $DJ(K)$  either of the two homotopy equivalent spaces.

**Corollary 3.8.** *The moment-angle complex  $\mathcal{Z}_K$  is the homotopy fibre of the embedding  $i: DJ(K) \hookrightarrow BT^m$ .*

**Corollary 3.9.** *The  $T^m$ -equivariant cohomology of  $\mathcal{Z}_K$  is isomorphic to the Stanley–Reisner ring of  $K$ :*

$$H_{T^m}^*(\mathcal{Z}_K) \cong \mathbb{Z}[K].$$

The following information about the homotopy groups of  $\mathcal{Z}_K$  can be retrieved from the above constructions.

**Proposition 3.10.** (a)  $\mathcal{Z}_K$  is 2-connected, and  $\pi_i(\mathcal{Z}_K) = \pi_i(DJ(K))$  for  $i \geq 3$ .

(b) If  $K$  is  $q$ -neighbourly, then  $\pi_i(\mathcal{Z}_K) = 0$  for  $i < 2q + 1$ . Moreover,  $\pi_{2q+1}(\mathcal{Z}_K)$  is a free Abelian group generated by the  $(q + 1)$ -element missing faces of  $K$ .

*Proof.* Note that  $BT^m = K(\mathbb{Z}^m, 2)$  and the 3-skeleton of  $DJ(K)$  coincides with that of  $BT^m$ . If  $K$  is  $q$ -neighbourly, then the  $(2q + 1)$ -skeleton of  $DJ(K)$  coincides with that of  $BT^m$ . Now, both statements follow easily from the exact homotopy sequence of the map  $i: DJ(K) \rightarrow BT^m$ .  $\square$

**3.2. Cell decompositions.** Here we construct a canonical cell decomposition of the moment-angle complex.

Let us decompose  $D^2$  into the union of one 2-cell  $D$ , one 1-cell  $T$  and one 0-cells 1, as shown on Figure 2. It defines a cellular complex structure on the polydisc  $(D^2)^m$  with  $3^m$  cells. Each cell of this complex is a product of cells of 3 different types:  $D_i$ ,  $T_i$  and  $1_i$ ,  $i = 1, \dots, m$ . We encode the cells using




FIGURE 2.

the language of “sign vectors”. Each cell of  $(D^2)^m$  will be represented by a sign vector  $\mathcal{T} \in \{D, T, 1\}^m$ . We denote by  $\mathcal{T}_D$ ,  $\mathcal{T}_T$  and  $\mathcal{T}_1$  respectively the  $D$ -,  $T$ - and  $1$ -component of  $\mathcal{T}$ . Each of these components can be seen as a subset of  $[m]$ , and all three subsets are complementary. The closure of a cell  $\mathcal{T}$  is homeomorphic to a product of  $|\mathcal{T}_D|$  discs and  $|\mathcal{T}_T|$  circles.

**Lemma 3.11.**  *$\mathcal{Z}_K$  is a cellular subcomplex of  $(D^2)^m$ . A cell  $\mathcal{T} \subset (D^2)^m$  belongs to  $\mathcal{Z}_K$  if and only if  $\mathcal{T}_D \in K$ .*

*Proof.* We have  $\mathcal{Z}_K = \cup_{\sigma \in K} B_\sigma$  and each  $B_\sigma$  is the closure of the cell  $\mathcal{T}$  with  $\mathcal{T}_D = \sigma$ ,  $\mathcal{T}_T = [m] \setminus \sigma$  and  $\mathcal{T}_1 = \emptyset$ .  $\square$

Denote by  $C^*(\mathcal{Z}_K)$  the corresponding cellular cochains. It has a natural bigrading defined by  $\text{bideg } \mathcal{T} = (-|\mathcal{T}_T|, 2|\mathcal{T}_T| + 2|\mathcal{T}_D|)$  (so  $\text{bideg } D_i = (0, 2)$ ,  $\text{bideg } T_i = (-1, 2)$  and  $\text{bideg } 1_i = (0, 0)$ ). Moreover, since the cellular differential does not change the second grading,  $C^*(\mathcal{Z}_K)$  splits into the sum of its components with fixed second degree:

$$C^*(\mathcal{Z}_K) = \bigoplus_{j=1}^m C^{*, 2j}(\mathcal{Z}_K).$$

Correspondingly, the cohomology of  $\mathcal{Z}_K$  acquires an additional grading, and one may define the *bigraded Betti numbers*  $b^{-i, 2j}(\mathcal{Z}_K)$  by

$$b^{-i, 2j}(\mathcal{Z}_K) := \dim_{\mathbf{K}} H^{-i, 2j}(\mathcal{Z}_K), \quad i, j = 1, \dots, m.$$

For the ordinary Betti numbers one has  $b^k(\mathcal{Z}_K) = \sum_{2j-i=k} b^{-i, 2j}(\mathcal{Z}_K)$ .

**Lemma 3.12.** *Let  $\varphi: K_1 \rightarrow K_2$  be a simplicial map between complexes on the sets  $[m_1]$  and  $[m_2]$  respectively. Then there is an equivariant cellular map  $\varphi_{\mathcal{Z}}: \mathcal{Z}_{K_1} \rightarrow \mathcal{Z}_{K_2}$  covering the induced map  $|\text{cone } K_1| \rightarrow |\text{cone } K_2|$ .*

*Proof.* Define the map  $\varphi_D: (D^2)^{m_1} \rightarrow (D^2)^{m_2}$  by

$$\varphi_D(z_1, \dots, z_{m_1}) = (w_1, \dots, w_{m_2}),$$

where

$$w_i = \prod_{j \in f^{-1}(i)} z_j.$$

Then one easily checks that  $\varphi_D(B_\sigma) \subset B_{\varphi(\sigma)}$ . Since  $\varphi$  is simplicial,  $\sigma \in K_1$  implies  $\varphi(\sigma) \in K_2$ . Hence, the restriction of  $\varphi_D$  to  $\mathcal{Z}_{K_1}$  is the required map.  $\square$



The above constructed map regards the bigrading, so the bigraded Betti numbers are functorial.

**3.3. Toric varieties, quasitoric manifolds, and torus manifolds.** Several important classes of manifolds with torus action emerge as the quotients of moment-angle complexes by appropriate freely acting subtori.

First we give the following characterisation of lsop's in the Stanley–Reisner ring. Let  $K^{n-1}$  be a simplicial complex and  $\theta_1, \dots, \theta_n$  a sequence of degree-two elements in  $\mathbf{k}[K]$ . We may write

$$(3.13) \quad \theta_i = \lambda_{i1}v_1 + \dots + \lambda_{im}v_m, \quad i = 1, \dots, n.$$

For arbitrary simplex  $\sigma \in K$  we have  $K_\sigma = \Delta^{|\sigma|-1}$  and  $\mathbf{k}[K_\sigma]$  is the polynomial ring  $\mathbf{k}[v_i : i \in \sigma]$  on  $|\sigma|$  generators. The inclusion  $K_\sigma \subset K$  induces the *restriction homomorphism*  $r_\sigma$  from  $\mathbf{k}[K]$  to the polynomial ring, mapping  $v_i$  identically if  $i \in \sigma$  and to zero otherwise. The degree-two part of a polynomial ring on  $q$  generators may be identified with the space of linear forms on  $\mathbf{k}^q$ .

**Lemma 3.14.** *A degree-two sequence  $\theta_1, \dots, \theta_n$  is an lsop in  $\mathbf{k}[K^{n-1}]$  if and only if its image under any restriction homomorphism  $r_\sigma$  generates  $(\mathbf{k}^{|\sigma|})^*$ .*

*Proof.* Suppose (3.13) is an lsop. For simplicity we denote its image under any restriction homomorphism by the same letters. Then the restriction induces a homomorphism of quotient rings:

$$\mathbf{k}[K]/(\theta_1, \dots, \theta_n) \rightarrow \mathbf{k}[v_i : i \in \sigma]/(\theta_1, \dots, \theta_n).$$

Since (3.13) is an lsop,  $\mathbf{k}[K]/(\theta_1, \dots, \theta_n)$  is a finitely generated  $\mathbf{k}$ -module. Hence, so is  $\mathbf{k}[v_i : i \in \sigma]/(\theta_1, \dots, \theta_n)$ . But the latter can be finitely generated only if  $\theta_1, \dots, \theta_n$  generates the degree-two part of the polynomial ring.

The “if” part may be proved by considering the sum of restrictions:

$$\mathbf{k}[K] \rightarrow \bigoplus_{\sigma \in K} \mathbf{k}[v_i : i \in \sigma],$$

which is actually a monomorphism. See [1, Thm. 5.1.16] for details.  $\square$

In particular, if  $K^{n-1}$  is pure (i.e., all maximal simplices have the same dimension), then (3.13) is an lsop if and only if its restriction to every  $(n-1)$ -simplex is a basis in the space of linear forms.

Suppose now that  $K$  is Cohen–Macaulay (e.g.,  $K$  is a sphere triangulation). Then every lsop is a regular sequence (however, for  $\mathbf{k} = \mathbb{Z}$  or a field of finite characteristic an lsop may fail to exist).

Now we restrict to the case  $\mathbf{k} = \mathbb{Z}$  and organise the coefficients in (3.13) into an  $n \times m$ -matrix  $\Lambda = (\lambda_{ij})$ . For arbitrary maximal simplex  $\sigma \in K$  denote by  $\Lambda_\sigma$  the square submatrix (minor) formed by the elements  $\lambda_{ij}$  with  $j \in \sigma$ . The matrix  $\Lambda$  defines a linear map  $\mathbb{Z}^m \rightarrow \mathbb{Z}^n$  and a homomorphism  $T^m \rightarrow T^n$ . We denote both by  $\lambda$  and denote the kernel of the latter map by  $T_\Lambda$ .

**Theorem 3.15.** *The following conditions are equivalent:*

- (a) *the sequence (3.13) is an lsop in  $\mathbb{Z}[K^{n-1}]$ ;*
- (b)  *$\det \Lambda_\sigma = \pm 1$  for every maximal simplex  $\sigma \in K$ ;*
- (c)  *$T_\Lambda \cong T^{m-n}$  and  $T_\Lambda$  acts freely on  $\mathcal{Z}_K$ .*

*Proof.* The equivalence of (a) and (b) is a reformulation of Lemma 3.14. Let us prove the equivalence of (b) and (c). Every isotropy subgroup of the  $T^m$ -action on  $\mathcal{Z}_K$  has the form

$$T^\sigma = \{(z_1, \dots, z_m) \in T^m : z_i = 1 \text{ if } i \notin \sigma\}$$

for some simplex  $\sigma \in K$ . Now, (b) is equivalent to the condition  $T_\Lambda \cap T^\sigma = \{e\}$  for arbitrary maximal  $\sigma$ , whence the statement follows.  $\square$

We denote the quotient  $\mathcal{Z}_K/T_\Lambda$  by  $M_K^{2n}(\Lambda)$  (when the context allows we may abbreviate it to  $M_K^{2n}$  or even to  $M^{2n}$ ). If  $K$  is a triangulated sphere, then  $\mathcal{Z}_K$  is a manifold, hence, so is  $M_K^{2n}$ . The  $n$ -torus  $T^n = T^m/T_\Lambda$  acts on  $M_K^{2n}$ . This construction generalises the following two important classes of  $T^n$ -manifolds.

Let  $K = K_P$  be a polytopal triangulation, dual to the boundary complex of a simple polytope  $P^n$ . Then the map  $\lambda$  determined by the matrix  $\Lambda$  may be regarded as an assignment of an integer vector to any facet of  $P^n$ . A map  $\lambda$  coming from a matrix satisfying the condition of Theorem 3.15(b) was called a *characteristic map* by Davis and Januszkiewicz. We refer to the corresponding quotient  $M_P^{2n}(\Lambda) = \mathcal{Z}_{K_P}/T_\Lambda$  as a *quasitoric manifold* (a toric manifold in the Davis–Januszkiewicz terminology).

Let us assume further that  $P^n$  is realised in  $\mathbb{R}^n$  with integer vertex coordinates, so we can write

$$(3.16) \quad P^n = \{\mathbf{x} \in \mathbb{R}^n : \langle \mathbf{l}_i, \mathbf{x} \rangle \geq -a_i, i = 1, \dots, m\},$$

where  $\mathbf{l}_i$  are integral vectors normal to the facets of  $P^n$  (we may assume these vectors to be primitive and inward pointing), and  $a_i \in \mathbb{Z}$ . Let  $\Lambda$  be the matrix formed by the column vectors  $\mathbf{l}_i$ ,  $i = 1, \dots, m$ . Then  $\mathcal{Z}_{K_P}/T_\Lambda$  is the *projective toric variety* determined by the polytope  $P^n$ . The condition of Theorem 3.15(b) is equivalent to that the toric variety is non-singular. Thereby a non-singular projective toric variety is a quasitoric manifold (but there are many quasitoric manifolds which are not toric varieties).

Finally, we mention that if  $K$  is an arbitrary triangulation of sphere, then the manifold  $M_K^{2n}(\Lambda)$  is a *torus manifold* in the sense of Hattori–Masuda [4]. The corresponding multi-fan has  $K$  as the underlying simplicial complex. This particular class of torus manifolds has many interesting specific properties.

#### 4. COHOMOLOGY OF MOMENT-ANGLE COMPLEXES

Studying ordinary and  $T^m$ -equivariant topology of  $\mathcal{Z}_K$  opens the way to some combinatorial applications. While the cohomology of  $DJ(K)$  is  $\mathbb{Z}[K]$ , the cohomology of the moment-angle complex  $\mathcal{Z}_K$  is isomorphic to the Tor-algebra  $\text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k})$ . The argument uses some algebraic topology techniques, such as Eilenberg–Moore spectral sequences (shortly *emss*). The intrinsic bigrading in Tor is exactly that coming from the bigraded cell decomposition of  $\mathcal{Z}_K$ . In the case when  $K$  is a triangulated manifold, the bigraded relative Poincaré duality for  $\mathcal{Z}_K$  gives the generalised Dehn–Sommerville equations.

**4.1. Eilenberg–Moore spectral sequence.** The Eilenberg–Moore spectral sequence can be considered as an extension of Adams’ cobar construction approach to calculating the cohomology of loop spaces. Here we give the necessary information on the spectral sequence; we follow L. Smith’s paper [5] in this description. The following theorem provides an algebraic setup for the emss.

**Theorem 4.1** (Eilenberg–Moore). *Let  $A$  be a differential graded  $\mathbf{k}$ -algebra, and  $M, N$  differential graded  $A$ -modules. Then there exists a second quadrant spectral sequence  $\{E_r, d_r\}$  converging to  $\mathrm{Tor}_A(M, N)$  whose  $E_2$ -term is*

$$E_2^{-i,j} = \mathrm{Tor}_{H[A]}^{-i,j}(H[M], H[N]), \quad i, j \geq 0,$$

where  $H[\cdot]$  denotes the cohomology algebra (or module).

Topological applications of the above theorem arise in the case when  $A, M, N$  are singular cochain algebras (or their commutative models) of certain topological spaces. The classical situation is described by the commutative diagram

$$(4.2) \quad \begin{array}{ccc} E & \longrightarrow & E_0 \\ \downarrow & & \downarrow \\ B & \longrightarrow & B_0, \end{array}$$

where  $E_0 \rightarrow B_0$  is a Serre fibre bundle with fibre  $F$  over a simply connected base  $B_0$ , and  $E \rightarrow B$  is the pullback along a continuous map  $B \rightarrow B_0$ . For any space  $X$ , let  $C^*(X; \mathbf{k})$  denote the singular cochain algebra of  $X$  or its commutative model (e.g., the cellular cochain algebra if a nice cellular product exists). Obviously,  $C^*(E_0; \mathbf{k})$  and  $C^*(B; \mathbf{k})$  are  $C^*(B_0; \mathbf{k})$ -modules. Under these assumptions the following statement holds.

**Lemma 4.3** (Eilenberg–Moore).  *$\mathrm{Tor}_{C^*(B_0; \mathbf{k})}(C^*(E_0; \mathbf{k}), C^*(B; \mathbf{k}))$  is an algebra in a natural way, and there is a canonical isomorphism of algebras*

$$\mathrm{Tor}_{C^*(B_0; \mathbf{k})}(C^*(E_0; \mathbf{k}), C^*(B; \mathbf{k})) \rightarrow H^*(E; \mathbf{k}).$$

The above two results lead to the following statement.

**Theorem 4.4** (Eilenberg–Moore). *There exists a spectral sequence of  $\mathbf{k}$ -algebras  $\{E_r, d_r\}$  with*

- (a)  $E_r \Rightarrow H^*(E; \mathbf{k})$ ;
- (b)  $E_2^{-i,j} = \mathrm{Tor}_{H^*(B_0; \mathbf{k})}^{-i,j}(H^*(E_0; \mathbf{k}), H^*(B; \mathbf{k}))$ .

The case when  $B$  above is a point is particularly important for applications, so we state the corresponding result separately.

**Corollary 4.5.** *Let  $E \rightarrow B$  be a fibration over a simply connected space  $B$  with fibre  $F$ . Then there exists a spectral sequence of  $\mathbf{k}$ -algebras  $\{E_r, d_r\}$  with*

- (a)  $E_r \Rightarrow H^*(F; \mathbf{k})$ ;
- (b)  $E_2 = \mathrm{Tor}_{H^*(B; \mathbf{k})}(H^*(E; \mathbf{k}), \mathbf{k})$ .

We refer to the above spectral sequence as the *Eilenberg–Moore spectral sequence of fibration  $E \rightarrow B$* .

Here is the first application of the emss. We use the notation of Theorem 3.15.

**Theorem 4.6.** *The cohomology of the quotient  $M_K(\Lambda) = \mathcal{Z}_K/T_\Lambda$  is given by*

$$H^*(M_K(\Lambda)) \cong \mathbb{Z}[K]/(\theta_1, \dots, \theta_n).$$

*Proof.* Denote  $M := M_K(\Lambda)$ . Since  $T_\Lambda$  acts freely on  $\mathcal{Z}_K$ , we have

$$ET^m \times_{T^m} \mathcal{Z}_K \simeq ET^n \times_{T^n} M,$$

where  $T^n = T^m/T_\Lambda$ . Hence,  $H^*(ET^n \times_{T^n} M) \cong \mathbb{Z}[K]$  by Corollary 3.9. The emss of the bundle  $ET^n \times_{T^n} M \rightarrow BT^n$  converges to  $H^*(M)$  and has

$$E_2^{*,*} = \mathrm{Tor}_{H^*(BT^n)}^{*,*}(H^*(ET^n \times_{T^n} M), \mathbb{Z}) = \mathrm{Tor}_{\mathbb{Z}[t_1, \dots, t_n]}^{*,*}(\mathbb{Z}[K], \mathbb{Z}),$$

where the  $\mathbb{Z}[t_1, \dots, t_n]$ -module structure in  $\mathbb{Z}[K]$  is defined through the homomorphism  $\mathbb{Z}[t_1, \dots, t_n] \rightarrow \mathbb{Z}[K]$  taking  $t_i$  to  $\theta_i$ ,  $i = 1, \dots, n$ , see (3.13). Since  $\mathbb{Z}[K]$  is Cohen–Macaulay and  $\theta_1, \dots, \theta_n$  is an lsop,  $\mathbb{Z}[K]$  is a free  $\mathbb{Z}[t_1, \dots, t_n]$ -module, so we have

$$\begin{aligned} \mathrm{Tor}_{\mathbb{Z}[t_1, \dots, t_n]}^{*,*}(\mathbb{Z}[K], \mathbb{Z}) &= \mathrm{Tor}_{\mathbb{Z}[t_1, \dots, t_n]}^{0,*}(\mathbb{Z}[K], \mathbb{Z}) \\ &= \mathbb{Z}[K] \otimes_{\mathbb{Z}[t_1, \dots, t_n]} \mathbb{Z} = \mathbb{Z}[K]/(\theta_1, \dots, \theta_n). \end{aligned}$$

Therefore,  $E_2^{0,*} = \mathbb{Z}[K]/(\theta_1, \dots, \theta_n)$  and  $E_2^{-p,*} = 0$  for  $p > 0$ . It follows that the Eilenberg–Moore spectral sequence collapses at the  $E_2$  term and  $H^*(M) = \mathbb{Z}[K]/(\theta_1, \dots, \theta_n)$ , as claimed.  $\square$

In the case of quasitoric manifold the cohomology ring was described in [3], while the case of non-singular toric varieties was known long before as the *Danilov–Jurkiewicz theorem*.

**Corollary 4.7.** *The cohomology of  $M_K(\Lambda)$  vanishes in odd dimensions and  $H^{2i}(M_K(\Lambda))$  is a free abelian group of rank  $h_i$ ,  $i = 0, \dots, n$ .*

In the case when  $P^n$  is an integral polytope (3.16) and  $M_P^{2n}$  a non-singular toric variety, the cohomology ring  $H^*(M_P)$  has more specific properties. Namely there is a cohomology class  $\omega \in H^2(M_P)$  for which the multiplication maps

$$H^{n-i}(M_P) \xrightarrow{\omega^i} H^{n+i}(M_P), \quad i = 1, \dots, n$$

are isomorphisms. This fact is known as the *Hard Lefschetz theorem for toric varieties* (although its proof is far beyond the scope of this review). One can take  $\omega = a_1 v_1 + \dots + a_m v_m$  in the notation of (3.16). The multiplication by  $\omega$  determines a monomorphism  $H^{2i}(M_P) \rightarrow H^{2i+2}(M_P)$  for  $i \leq \lfloor \frac{n}{2} \rfloor$ . Consider the graded algebra  $A := H^*(M_P)/(\omega)$ . It is generated in degree two and has  $\dim A^{2i} = h_i - h_{i-1}$ ,  $i = 1, \dots, \lfloor \frac{n}{2} \rfloor$ . Then it follows from Theorem 1.7 that

$$0 \leq h_{i+1} - h_i \leq (h_i - h_{i-1})^{(i)}$$

for  $i < \lfloor \frac{n}{2} \rfloor$ , which proves the necessity part of Theorem 1.3.

**4.2. Cohomology of  $\mathcal{Z}_K$ .** Here we calculate the cohomology algebra of  $\mathcal{Z}_K$ . As an immediate corollary we obtain that the cohomology inherits a canonical bigrading from the spectral sequence. The corresponding bigraded Betti numbers coincide with the algebraic Betti numbers of  $\mathbf{k}[K]$  introduced before (2.22).

**Theorem 4.8.** *There are isomorphisms of algebras*

$$H^*(\mathcal{Z}_K; \mathbf{k}) \cong \mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k}) \cong H[\mathbf{k}[K] \otimes \Lambda[u_1, \dots, u_m], d],$$

where the latter algebra is bigraded by  $\mathrm{bideg} v_i = (0, 2)$ ,  $\mathrm{bideg} u_i = (-1, 2)$  and the differential is determined by  $dv_i = 0$ ,  $du_i = v_i$ . In particular,

$$\dim_{\mathbf{k}} H^q(\mathcal{Z}_K; \mathbf{k}) \cong \sum_{-i+2j=q} \beta^{-i, 2j}(\mathbf{k}(K)).$$

*Proof.* Consider the emss of the commutative square

$$\begin{array}{ccc} E & \longrightarrow & ET^m \\ \downarrow & & \downarrow \\ DJ(K) & \xrightarrow{i} & BT^m \end{array},$$

where the left vertical arrow is the pullback along  $i$ . Corollary 3.8 shows that  $E$  is homotopy equivalent to  $\mathcal{Z}_K$ .

By Proposition 3.1, the map  $i: DJ(K) \hookrightarrow BT^m$  induces the quotient epimorphism  $i^*: \mathbf{k}[v_1, \dots, v_m] \rightarrow \mathbf{k}[K]$  of the cellular cochain algebras. Since  $ET^m$  is contractible, there is a chain equivalence  $C^*(ET^m; \mathbf{k}) \simeq \mathbf{k}$ . More precisely,  $C^*(ET^m; \mathbf{k})$  can be identified with the Koszul resolution  $\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[v_1, \dots, v_m]$  of  $\mathbf{k}$  (see Example 2.19). Therefore, we have an isomorphism

$$\mathrm{Tor}_{C^*(BT^m)}(C^*(DJ(K)), C^*(ET^m)) \cong \mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k}).$$

On the other hand, Lemma 4.3 shows that  $\mathrm{Tor}_{C^*(BT^m)}(C^*(DJ(K)), C^*(ET^m))$  is an algebra isomorphic to  $H^*(\mathcal{Z}_K; \mathbf{k})$ , which proves the first isomorphism. The second one follows from Lemma 2.26.  $\square$

**Exercise 4.9.** Prove that the Leray–Serre spectral sequence of the principal  $T^m$ -bundle  $\mathcal{Z}_K \rightarrow DJ(K)$  collapses at the  $E_3$  term.

**Example 4.10.** Let  $K = \partial\Delta^{m-1}$ . Then  $\mathbf{k}[K] = \mathbf{k}[v_1, \dots, v_m]/(v_1 \cdots v_m)$ . A direct calculation shows that the cohomology of  $\mathbf{k}[K] \otimes \Lambda[u_1, \dots, u_m]$  is additively generated by the classes 1 and  $[v_1 v_2 \cdots v_{m-1} u_m]$ . The latter represents the fundamental cohomology class of  $\mathcal{Z}_K \cong S^{2m-1}$ .

**Example 4.11.** Let  $K$  be the boundary complex of an  $m$ -gon  $P^2$  with  $m \geq 4$ . We have  $\mathbf{k}[K] = \mathbf{k}[v_1, \dots, v_m]/\mathcal{I}_K$ , where  $\mathcal{I}_K$  is generated by the monomials  $v_i v_j$ ,  $i - j \neq 0, 1 \pmod m$ . The complex  $\mathcal{Z}_K$  is a manifold of dimension  $m + 2$ . For  $m = 4$  we have  $\mathcal{Z}_K = S^3 \times S^3$ . Suppose  $m = 5$ . Then the group  $H^3(\mathcal{Z}_K)$  has 5 generators represented by the cocycles  $v_i u_{i+2} \in \mathbf{k}[K] \otimes \Lambda[u_1, \dots, u_5]$ ,  $i = 1, \dots, 5$ , and  $H^4(\mathcal{Z}_K)$  has 5 generators represented by the cocycles  $v_j u_{j+2} u_{j+3}$ ,  $j = 1, \dots, 5$ . The product of  $v_i u_{i+2}$  and  $v_j u_{j+2} u_{j+3}$  represents a non-zero cohomology class in  $H^7(\mathcal{Z}_K)$  if and only if all the indices  $i, i + 2, j, j + 2, j + 3$  are different. Thus, for each of the 5 cohomology classes  $[v_i u_{i+2}]$  there is a unique (Poincaré dual) cohomology class  $[v_j u_{j+2} u_{j+3}]$  such that the product

$[v_i u_{i+2}] \cdot [v_j u_{j+2} u_{j+3}]$  is non-zero. This describes the multiplicative structure in the cohomology of  $\mathcal{Z}_K$ . In particular, its Betti vector is  $(1, 0, 0, 5, 5, 0, 0, 1)$ . There is also a similar description for  $m > 5$ , see [2, Ex. 7.22].

Now we are going to compare the bigrading in  $H^*(\mathcal{Z}_K)$  determined by Theorem 4.8 with that coming from the bigraded cellular structure.

Define the map

$$\begin{aligned} j: \mathbf{k}[K] \otimes \Lambda[u_1, \dots, u_m] &\rightarrow C^*(\mathcal{Z}_K; \mathbf{k}), \\ v_i &\mapsto D_i^*, \quad u_i \mapsto T_i^*. \end{aligned}$$

Hence,  $j$  maps a monomial  $v_\sigma u_\tau$  with  $\sigma \cap \tau = \emptyset$  to the cellular cochain  $\mathcal{T}(\sigma, \tau)^*$  with  $\mathcal{T}(\sigma, \tau)_D = \sigma$ ,  $\mathcal{T}(\sigma, \tau)_T = \tau$  (note that  $\sigma \in K$ ) and all other monomials to zero. The above map respects the bigrading in both algebras, and one directly checks that it commutes with the differentials.

**Theorem 4.12.** *The map  $j$  is a quasiisomorphism, that is, induces an isomorphism  $\mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k}) \rightarrow H^*(\mathcal{Z}_K; \mathbf{k})$  in the cohomology.*

*Proof.* The map  $j$  has an obvious additive right inverse  $i: C^*(\mathcal{Z}_K; \mathbf{k}) \rightarrow \mathbf{k}[K] \otimes \Lambda[u_1, \dots, u_m]$  sending  $T(\sigma, \tau)$  to  $v_\sigma u_\tau$ . The standard cochain homotopy operator  $s$  for the Koszul resolution establishes a cochain homotopy equivalence between  $\mathrm{id}$  and  $ij$ , that is,  $ds + sd = \mathrm{id} - ij$ . See [2, Ch. 7] for more details.  $\square$

**Corollary 4.13.** *The algebraic bigraded Betti numbers of  $\mathbf{k}[K]$  coincide with the topological bigraded Betti numbers of  $\mathcal{Z}_K$ :*

$$\beta^{-i, 2j}(\mathbf{k}[K]) = b^{-i, 2j}(\mathcal{Z}_K; \mathbf{k}), \quad i, j = 1, \dots, m.$$

Now we can summarise the results of Proposition 2.3, Lemmas 2.27 and 3.12, Corollary 3.9 and Theorem 4.12 in a statement describing the functorial properties of the correspondence  $K \mapsto \mathcal{Z}_K$ . Let us introduce the following functors:

- $\mathcal{Z}$ , the covariant functor  $K \mapsto \mathcal{Z}_K$  from the category of finite simplicial complexes and simplicial maps to the category of toric spaces and equivariant maps (the *moment-angle complex functor*);
- $\mathbf{k}[\cdot]$ , the contravariant functor  $K \mapsto \mathbf{k}[K]$  from simplicial complexes to graded  $\mathbf{k}$ -algebras (the *Stanley–Reisner functor*);
- $\mathrm{Tor}\text{-alg}$ , the contravariant functor

$$K \mapsto \mathrm{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k})$$

from complexes to bigraded  $\mathbf{k}$ -algebras (the *Tor-algebra functor*);

- $H_T^*$ , the contravariant functor  $X \mapsto H_T^*(X; \mathbf{k})$  from toric spaces and equivariant maps to  $\mathbf{k}$ -algebras (the *equivariant cohomology functor*);
- $H^*$ , the contravariant functor  $X \mapsto H^*(X; \mathbf{k})$  from spaces to  $\mathbf{k}$ -algebras (the *ordinary cohomology functor*).

**Proposition 4.14.** *We have the following identities of functors:*

$$H_T^* \circ \mathcal{Z} = \mathbf{k}[\cdot], \quad H^* \circ \mathcal{Z} = \mathrm{Tor}\text{-alg}.$$

The later identity means that for every simplicial map  $\varphi: K_1 \rightarrow K_2$  the cohomology map  $\varphi_{\mathcal{Z}}^*: H^*(\mathcal{Z}_{K_2}; \mathbf{k}) \rightarrow H^*(\mathcal{Z}_{K_1}; \mathbf{k})$  coincides with the induced homomorphism  $\varphi_t^*$  (2.28) of Tor-algebras.

Also, now we are ready to give a proof of Theorem 2.24.

*Proof of the Hochster theorem.* First we observe that the cellular cochains of  $\mathcal{Z}_K$  decompose as

$$(4.15) \quad C^{*,*}(\mathcal{Z}_K) = \bigoplus_{\omega \subset [m]} C^{*,2\omega}(\mathcal{Z}_K)$$

(as bigraded differential modules), where  $C^{*,2\omega}(\mathcal{Z}_K)$  is the subcomplex generated by the cochains  $\mathcal{T}(\sigma, \omega \setminus \sigma)^*$  with  $\sigma \subset \omega$  (remember that  $\mathcal{T}(\sigma, \omega \setminus \sigma)_D = \sigma$  and  $\mathcal{T}(\sigma, \omega \setminus \sigma)_T = \omega \setminus \sigma$ ). It follows that

$$(4.16) \quad b^{-i,2j}(\mathcal{Z}_K) = \sum_{\omega \subset [m]: |\omega|=j} b^{-i,2\omega}(\mathcal{Z}_K),$$

where  $b^{-i,2\omega}(\mathcal{Z}_K) := \dim H^{-i}[C^{*,2\omega}(\mathcal{Z}_K)]$ .

The assignment  $\sigma^* \mapsto \mathcal{T}(\sigma, \omega \setminus \sigma)^* \subset C^{|\sigma|-|\omega|,2\omega}(\mathcal{Z}_K)$  defines an isomorphism of cochain complexes

$$C^*(K_\omega) \rightarrow C^{*+1-|\omega|,2\omega}(\mathcal{Z}_K),$$

where the former is the simplicial cochains of  $K_\omega$ . Hence,

$$b^{-i,2\omega}(\mathcal{Z}_K) = b^{|\omega|-i-1}(K_\omega).$$

This together with (4.16) and Corollary 4.13 implies the Hochster theorem.  $\square$

**4.3. Generalised Dehn–Sommerville equations.** Here we consider the bigraded Poincaré duality for  $\mathcal{Z}_K$ . As a corollary, we deduce linear relations for the number of faces in a triangulated manifold, generalising the Dehn–Sommerville equations.

The decomposition (4.15) gives rise to the coarser decomposition

$$C^{*,*}(\mathcal{Z}_K) = \bigoplus_{p=0}^m C^{*,2p}(\mathcal{Z}_K).$$

Let us consider the corresponding Euler characteristics:

$$\chi_p(\mathcal{Z}_K) := \sum_{q=0}^m (-1)^q \dim C^{-q,2p}(\mathcal{Z}_K) = \sum_{q=0}^m (-1)^q b^{-q,2p}(\mathcal{Z}_K)$$

and define the generating polynomial  $\chi(\mathcal{Z}_K; t)$  by

$$\chi(\mathcal{Z}_K; t) = \sum_{p=0}^m \chi_p(\mathcal{Z}_K) t^{2p}.$$

It turns out that this polynomial can be expressed in terms of the number of faces of  $K$ . Introduce the *h-polynomial* of  $K$  as  $h(t) = h_0 + h_1 t + \cdots + h_n t^n$ , where  $(h_0, h_1, \dots, h_n)$  is the *h-vector*.

**Lemma 4.17.** *For every  $K^{n-1}$  it holds that*

$$\chi(\mathcal{Z}_K; t) = (1 - t^2)^{m-n} h(t^2).$$

*Proof.* The component  $C^{-q, 2p}(\mathcal{Z}_K)$  has basis of cochains  $\mathcal{T}(\sigma, \tau)^*$  with  $\sigma \in K$ ,  $|\sigma| = p - q$  and  $|\tau| = q$ . It follows that

$$\dim C^{-q, 2p}(\mathcal{Z}_K) = f_{p-q-1} \binom{m-p+q}{q}.$$

The rest is a direct calculation using (1.1). See [2, Thm. 7.15] for details.  $\square$

We may regard the standard torus  $T^m \subset \mathbb{C}^m$  as a cellular subcomplex in the moment-angle complex  $\mathcal{Z}_K$ . In the same fashion as we did before for  $\mathcal{Z}_K$ , we may define the bigraded Betti numbers and characteristic polynomials for the pair  $(\mathcal{Z}_K, T^m)$  and for  $\mathcal{Z}_K \setminus T^m$ . The proof of the following statement uses a similar but rather more complicated argument as that of Lemma 4.17.

**Lemma 4.18.** *For every  $K^{n-1}$  it holds that*

$$\begin{aligned} \chi(\mathcal{Z}_K, T^m; t) &= (1 - t^2)^{m-n} h(t^2) - (1 - t^2)^m; \\ \chi(\mathcal{Z}_K \setminus T^m; t) &= (1 - t^2)^{m-n} h(t^2) + (-1)^{n-1} h_n (1 - t^2)^m. \end{aligned}$$

Assume that  $K$  is a triangulation of  $S^{n-1}$ . Then  $\mathcal{Z}_K$  is an  $(m+n)$ -dimensional (closed) manifold.

**Exercise 4.19.** Show that the top cohomology group  $H^{m+n}(\mathcal{Z}_K)$  is generated by the cohomology class of any monomial  $v_\sigma u_\tau \in \mathbf{k}[K] \otimes \Lambda[u_1, \dots, u_m]$  of bidegree  $-(m-n, 2m)$  satisfying  $\sigma \cap \tau = \emptyset$ .

**Corollary 4.20.** *The Poincaré duality for  $\mathcal{Z}_K$  respects the bigraded structure in the cohomology. In particular,*

$$b^{-q, 2p}(\mathcal{Z}_K) = b^{-(m-n)+q, 2(m-p)}(\mathcal{Z}_K).$$

It follows that

$$\chi_p(\mathcal{Z}_K) = (-1)^{m-n} \chi_{m-p}(\mathcal{Z}_K) \quad \text{and} \quad \chi(\mathcal{Z}_K; t) = (-1)^{m-n} t^{2m} \chi(\mathcal{Z}_K; \frac{1}{t}).$$

**Exercise 4.21.** Deduce the Dehn–Sommerville equations  $h_i = h_{n-i}$  from the above identity and Lemma 4.17.

Now assume that  $K$  is a triangulation of a closed  $(n-1)$ -dimensional manifold. In this case  $\mathcal{Z}_K$  may fail to be a manifold, but  $\mathcal{Z}_K \setminus T^m$  still has a homotopy type of a manifold with boundary. The relative Poincaré duality still regards the bigradings, so we have

$$(4.22) \quad \chi(\mathcal{Z}_K \setminus T^m; t) = (-1)^{m-n} t^{2m} \chi(\mathcal{Z}_K, T^m; \frac{1}{t}).$$


**Theorem 4.23** (Dehn–Sommerville equations for triangulated manifolds). *The following relations hold for the  $h$ -vector  $(h_0, h_1, \dots, h_n)$  of any triangulated manifold  $K^{n-1}$ :*

$$h_{n-i} - h_i = (-1)^i (\chi(K^{n-1}) - \chi(S^{n-1})) \binom{n}{i}, \quad i = 0, 1, \dots, n.$$

*Proof.* It follows from Lemma 4.18 and (4.22). See [2, Thm. 7.44] for details.  $\square$

Note that  $\chi(K^{n-1}) - \chi(S^{n-1}) = (-1)^{n-1} (h_n - 1)$ , so the above relations are indeed linear equations for the numbers of faces.





(a)  $\mathbf{f} = (9, 27, 18)$ ,  $\mathbf{h} = (1, 6, 12, -1)$       (b)  $\mathbf{f} = (7, 21, 14)$ ,  $\mathbf{h} = (1, 4, 10, -1)$

FIGURE 3. “Symmetric” and “minimal” triangulation of  $T^2$

**Example 4.24.** Consider triangulations of the 2-torus  $T^2$ . We have  $n = 3$ ,  $\chi(T^2) = 0$ . From  $\chi(K^{n-1}) = 1 + (-1)^{n-1}h_n$  we deduce  $h_3 = -1$ . The Dehn–Sommerville equations are

$$h_3 - h_0 = -2, \quad h_2 - h_1 = 6.$$

For instance, the triangulation on Figure 3 (a) has  $f_0 = 9$  vertices,  $f_1 = 27$  edges and  $f_2 = 18$  triangles. The corresponding  $h$ -vector is  $(1, 6, 12, -1)$ .

On the other hand, a triangulation of  $T^2$  with only 7 vertices can be achieved, see Figure 3 (b). Note that this triangulation is neighbourly, i.e. its 1-skeleton is a complete graph on 7 vertices.

**Exercise 4.25.** Show that any triangulation of  $T^2$  has at least 7 vertices.

## 5. COORDINATE SUBSPACE ARRANGEMENTS

Yet another construction of the moment-angle complex  $\mathcal{Z}_K$  comes from the study of coordinate subspace arrangements. Coordinate subspace arrangements in  $m$ -dimensional complex space are in functorial one-to-one correspondence with simplicial complexes on  $m$  vertices. The complement to a coordinate subspace arrangement is homotopy equivalent to moment-angle complex  $\mathcal{Z}_K$  for the corresponding simplicial complex  $K$ . We investigate different consequences of this fact. In particular, we compare the Goresky–MacPherson calculations of Betti numbers of a coordinate subspace arrangement complement with Hochster’s calculation for the bigraded Betti numbers of the Tor-algebra.

A *complex coordinate subspace arrangement* is a set  $\mathcal{CA} = \{L_1, \dots, L_r\}$  of coordinate subspaces in some  $\mathbb{C}^m$ . Each coordinate subspace can be written as

$$(5.1) \quad L_\omega = \{(z_1, \dots, z_m) \in \mathbb{C}^m : z_{i_1} = \dots = z_{i_k} = 0\},$$

where  $\omega = \{i_1, \dots, i_k\}$  is a subset of  $[m]$ .

For each simplicial complex  $K$  on the set  $[m]$  we define the complex coordinate subspace arrangement  $\mathcal{CA}(K) := \{L_\omega : \omega \notin K\}$  and its *complement*

$$U(K) := \mathbb{C}^m \setminus \bigcup_{\omega \notin K} L_\omega.$$

Note that if  $L \subset K$  is a subcomplex, then  $U(L) \subset U(K)$ . The following observation is straightforward.

**Proposition 5.2.** *The assignment  $K \mapsto U(K)$  defines a one-to-one order-preserving correspondence between simplicial complexes on  $[m]$  and coordinate subspace arrangement complements in  $\mathbb{C}^m$ .*

**Example 5.3.** 1. If  $K = \Delta^{m-1}$  then  $U(K) = \mathbb{C}^m$ .

2. If  $K = \partial\Delta^{m-1}$  then  $U(K) = \mathbb{C}^m \setminus \{0\}$ .

3. If  $K$  is a disjoint union of  $m$  vertices, then  $U(K)$  is the complement in  $\mathbb{C}^m$  to the set of all codimension-two coordinate subspaces  $z_i = z_j = 0$ ,  $1 \leq i < j \leq m$ .

The complement  $U(K)$  is invariant under the diagonal action of  $T^m$  on  $\mathbb{C}^m$ .

**Proposition 5.4.** *There is an equivariant deformation retraction  $U(K) \rightarrow \mathcal{Z}_K$ .*

*Proof.* For arbitrary subset  $\omega \subset [m]$ , set

$$U_\omega := \{(z_1, \dots, z_m) \in \mathbb{C}^m : |z_i| \neq 0 \text{ if } i \notin \omega\}.$$

Then  $U(K) = \bigcup_{\sigma \in K} U_\sigma$ . Each  $B_\sigma$  in (3.2) is a deformation retract of  $U_\sigma$ , and these deformation retractions patch together to yield a deformation retraction  $U(K) \rightarrow \mathcal{Z}_K$ .  $\square$

Hence, we may use the previous results on moment-angle complexes to calculate the cohomology ring of a coordinate subspace arrangement complement.

**Corollary 5.5.** *The following isomorphism of graded algebras holds:*

$$H^*(U(K); \mathbf{k}) \cong \text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[K], \mathbf{k}) = H[\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}(K), d].$$

**Example 5.6.** Let  $K$  and  $U(K)$  be as in Example 5.3.3. Then  $\mathbf{k}(K) = \mathbf{k}[v_1, \dots, v_m]/\mathcal{I}_K$ , where  $\mathcal{I}_K$  is generated by the monomials  $v_i v_j$ ,  $i \neq j$ . An easy calculation using Corollary 5.5 shows that monomials  $v_{i_1} u_{i_2} u_{i_3} \cdots u_{i_k}$  with  $k \geq 2$  and  $i_p \neq i_q$  for  $p \neq q$  form a basis of cocycles in  $\mathbf{k}(K) \otimes \Lambda[u_1, \dots, u_m]$ . Since  $\deg(v_{i_1} u_{i_2} u_{i_3} \cdots u_{i_k}) = k + 1$ , the space of degree- $(k + 1)$  cocycles has dimension  $m \binom{m-1}{k-1}$ . The space of degree- $(k + 1)$  coboundaries is  $\binom{m}{k}$ -dimensional and is spanned by the coboundaries of the form  $d(u_{i_1} \cdots u_{i_k})$ . Hence,

$$\begin{aligned} \dim H^0(U(K)) &= 1, \quad H^1(U(K)) = H^2(U(K)) = 0, \\ \dim H^{k+1}(U(K)) &= m \binom{m-1}{k-1} - \binom{m}{k} = (k-1) \binom{m}{k}, \quad 2 \leq k \leq m, \end{aligned}$$

and the multiplication in the cohomology is trivial.

In particular, for  $m = 3$  we have 6 three-dimensional cohomology classes  $[v_i u_j]$ ,  $i \neq j$ , subject to 3 relations  $[v_i u_j] = [v_j u_i]$ , and 3 four-dimensional cohomology classes  $[v_1 u_2 u_3]$ ,  $[v_2 u_1 u_3]$ ,  $[v_3 u_1 u_2]$  subject to one relation

$$[v_1 u_2 u_3] - [v_2 u_1 u_3] + [v_3 u_1 u_2] = 0.$$

Hence,  $\dim H^3(U(K)) = 3$ ,  $\dim H^4(U(K)) = 2$ , and the multiplication is trivial. It can be shown that  $U(K)$  in this case has a homotopy type of a wedge of spheres:

$$U(K) \simeq S^3 \vee S^3 \vee S^3 \vee S^4 \vee S^4.$$

One can define a coordinate subspace as the linear span of a subset of the standard basis in  $\mathbb{C}^m$ , rather than by setting some coordinates to be zero as in (5.1). This gives an alternative way to parametrise coordinate subspace arrangements by simplicial complexes. Namely, we can write

$$\mathcal{CA}(K) = \{L_\omega : \omega \notin K\} = \{\text{span}\langle e_{i_1}, \dots, e_{i_k} \rangle : \{i_1, \dots, i_k\} \in \widehat{K}\}$$

where  $\widehat{K}$  is the simplicial complex given by

$$\widehat{K} := \{\omega \subset [m] : [m] \setminus \omega \notin K\}.$$

It is called the *dual complex* of  $K$ . The cohomology of full subcomplexes in  $K$  is related to the homology of links in  $\widehat{K}$  by means of the following combinatorial version of the Alexander duality theorem.

**Theorem 5.7** (Alexander duality). *For any simplicial complex  $K \neq \Delta^{m-1}$  on  $[m]$  and simplex  $\sigma \in \widehat{K}$  it holds that*

$$\widetilde{H}^i(K_{\widehat{\sigma}}) \cong \widetilde{H}_{m-3-i-|\sigma|}(\text{link}_{\widehat{K}} \sigma),$$

where  $\widehat{\sigma} := [m] \setminus \sigma$ . In particular, for  $\sigma = \emptyset$  we get

$$\widetilde{H}^i(K) \cong \widetilde{H}_{m-3-i}(\widehat{K}), \quad -1 \leq j \leq m-2,$$

The Alexander duality and Hochster theorem allow us to give a simple argument for the *Goresky–MacPherson formula* in the coordinate subspace arrangement case, providing another way to calculate the cohomology of the complement. The theorem below was firstly proved using elaborated algebraic geometry techniques, such as *stratified Morse theory* and *intersection cohomology*.

**Theorem 5.8** (Goresky–MacPherson). *We have*

$$\widetilde{H}^i(U(K)) = \bigoplus_{\sigma \in \widehat{K}} \widetilde{H}_{2m-2|\sigma|-i-2}(\text{link}_{\widehat{K}} \sigma).$$

*Proof.* Using Corollary 5.5 to identify  $\beta^{-i,2j}(\mathbf{k}[K])$  with  $\dim_{\mathbf{k}} H^{-i,2j}(U(K))$ , we get the following formula from Hochster’s Theorem 2.24:

$$H^p(U(K)) = \bigoplus_{\omega \subset [m]} \widetilde{H}^{p-|\omega|-1}(K_\omega).$$

Non-empty simplices  $\omega \in K$  do not contribute to the above sum, since the corresponding subcomplexes  $K_\omega$  are contractible. Since  $\widetilde{H}^{-1}(\emptyset) = \mathbf{k}$ , the empty subset of  $[m]$  only contributes  $\mathbf{k}$  to  $H^0(U(K))$ . Hence, we may rewrite the above formula as

$$\widetilde{H}^p(U(K)) = \bigoplus_{\omega \notin K} \widetilde{H}^{p-|\omega|-1}(K_\omega) = \bigoplus_{\sigma \in \widehat{K}} \widetilde{H}^{p-m+|\sigma|-1}(K_{\widehat{\sigma}}).$$

Using Theorem 5.7, we identify the latter with  $\bigoplus_{\sigma \in \widehat{K}} \widetilde{H}_{2m-2|\sigma|-p-2}(\text{link}_{\widehat{K}} \sigma)$ , as claimed.  $\square$

**Exercise 5.9.** Calculate the cohomology of  $U(K)$  from Example 5.6 using the Goresky–MacPherson formula.

## REFERENCES

- [1] Winfried Bruns and Jürgen Herzog, *Cohen-Macaulay rings*, revised edition, Cambridge Studies in Adv. Math. **39**, Cambridge Univ. Press, Cambridge, 1998.
- [2] Victor M. Buchstaber, Taras E. Panov, *Torus actions and their applications in topology and combinatorics*, University Lecture Series **24**, Amer. Math. Soc., Providence, R.I., 2002.
- [3] Michael W. Davis and Tadeusz Januszkiewicz, *Convex polytopes, Coxeter orbifolds and torus actions*, Duke Math. J. **62** (1991), no. 2, 417–451.
- [4] Akio Hattori and Mikiya Masuda, *Theory of multi-fans*, preprint (2001), arXiv: math.SG/0106229.
- [5] Larry Smith, *Homological algebra and the Eilenberg–Moore spectral sequence*, Trans. Amer. Math. Soc. **129** (1967), 58–93.
- [6] Richard P. Stanley, *Combinatorics and Commutative Algebra*, second edition, Progress in Math. **41**, Birkhäuser, Boston, 1996.

DEPARTMENT OF MATHEMATICS AND MECHANICS, MOSCOW STATE UNIVERSITY, MOSCOW 119992, RUSSIA

and INSTITUTE FOR THEORETICAL AND EXPERIMENTAL PHYSICS, MOSCOW 117259, RUSSIA

*E-mail address:* `tpanov@mech.math.msu.su`